



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

A Genome-wide Association Study of Emphysema and Airway Quantitative Imaging Phenotypes

Citation for published version:

NETT Genetics, ECLIPSE, and COPDGene Investigators, Cho, MH, Castaldi, PJ, Hersh, CP, Hobbs, BD, Barr, RG, Tal-Singer, R, Bakke, P, Gulsvik, A, San José Estépar, R, Van Beek, EJR, Coxson, HO, Lynch, DA, Washko, GR, Laird, NM, Crapo, JD, Beaty, TH & Silverman, EK 2015, 'A Genome-wide Association Study of Emphysema and Airway Quantitative Imaging Phenotypes', *American Journal of Respiratory and Critical Care Medicine*, vol. 192, no. 5, pp. 559-569. <https://doi.org/10.1164/rccm.201501-0148OC>

Digital Object Identifier (DOI):

[10.1164/rccm.201501-0148OC](https://doi.org/10.1164/rccm.201501-0148OC)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

American Journal of Respiratory and Critical Care Medicine

Publisher Rights Statement:

This is the author's accepted manuscript.

The final published version is available at: <http://www.atsjournals.org/doi/abs/10.1164/rccm.201501-0148OC#.VXWCLGPky30>

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Title**A Genome-wide Association Study of Emphysema and Airway Quantitative Imaging Phenotypes**

Michael H. Cho^{1,2}, Peter J. Castaldi¹, Craig P. Hersh^{1,2}, Brian D. Hobbs^{1,2}, R. Graham Barr⁴, Ruth Tal-Singer⁵, Per Bakke⁶, Amund Gulsvik⁶, Raúl San José Estépar³, Edwin Van Beek⁷, Harvey O. Coxson⁸, David A. Lynch⁹, George R. Washko², Nan M. Laird¹⁰, James D. Crapo⁹, Terri H. Beaty¹¹, Edwin K. Silverman^{1,2}, on behalf of the NETT Genetics, ECLIPSE, and COPDGene Investigators

¹Channing Division of Network Medicine, ²Division of Pulmonary and Critical Care Medicine, and ³Laboratory of Mathematics in Imaging, Department of Radiology, Brigham and Women's Hospital, Boston, MA; ⁴Department of Medicine, College of Physicians and Surgeons, and Department of Epidemiology, Mailman School of Public Health, Columbia University, New York, New York, ⁵GlaxoSmithKline Research and Development, King Of Prussia, PA, ⁶Department of Clinical Science, University of Bergen, Norway, ⁷Queens Medical Research Institute, University of Edinburgh, Department of Radiology and Department of Biomedical Engineering, University of Iowa; ⁸Department of Radiology, University of British Columbia, Vancouver, Canada ⁹National Jewish Health, Denver, CO; ¹⁰Harvard School of Public Health, Boston, MA; ¹¹Johns Hopkins University Bloomberg School of Public Health, Baltimore, MD.

Corresponding author

Michael H. Cho (remhc@channing.harvard.edu), tel: 617-525-0897, fax: 888-487-1078.

Author Contributions: Study design: MHC, CPH, RGB, RTS, PB, AG, HOC, DAL, GRW, NL, JDC, THB, EKS. Phenotype acquisition and quality control: MHC, PJC, CPH, RGB, RTS, PB, AG, HOC, DAL, GRW, JDC, EKS. Genotype data acquisition and quality control: MHC, PJC, THB, EKS. Data analysis: MHC, PJC, BDH, NML, THB, EKS. Critical revision of the manuscript: all authors

Funding:

This work was supported by NHLBI R01 HL084323, P01 HL083069, P01 HL105339 and R01 HL089856 (E.K.S.); K08 HL097029 and R01 HL113264 (M.H.C.), and R01 HL089897 (J.D.C.); and the Alpha-1 Foundation (M.H.C.). The COPDGene study (NCT00608764) is also supported by the COPD Foundation through contributions made to an Industry Advisory Board comprised of AstraZeneca, Boehringer Ingelheim, Novartis, Pfizer, GlaxoSmithKline, Siemens and Sunovion. The National Emphysema Treatment Trial was supported by the NHLBI N01HR76101, N01HR76102, N01HR76103, N01HR76104, N01HR76105, N01HR76106, N01HR76107, N01HR76108, N01HR76109, N01HR76110, N01HR76111, N01HR76112, N01HR76113, N01HR76114, N01HR76115, N01HR76116, N01HR76118 and N01HR76119, the Centers for Medicare and Medicaid Services and the Agency for Healthcare Research and Quality. The Norway GenKOLS study (Genetics of Chronic Obstructive Lung Disease, GSK code RES11080), the ECLIPSE study (NCT00292552; GSK code SCO104960) were funded by GlaxoSmithKline.

The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Heart, Lung, and Blood Institute or the National Institutes of Health.

Running Head: GWAS of Quantitative Imaging Phenotypes

Descriptor number: 9.6 COPD: Epidemiology

Word count: 3288

At a Glance Commentary

Scientific Knowledge on the Subject

Chronic obstructive pulmonary disease is a complex and heterogeneous disease. Quantitative image analysis of chest CT scans can characterize this heterogeneity. Recent studies have identified genetic variants that increase susceptibility to emphysema or airway wall thickening, but have not examined both measurements in large populations of subjects with disease.

What This Study Adds to the Field

Our study confirms previously described associations and additionally identifies new genome-wide significant associations with emphysema near *SERPINA10* and *DLC1*. We also show that many loci previously identified in population-based studies of lung function are associated with emphysema or airway phenotypes. Genome-wide analysis of quantitative imaging may identify novel risk factors for COPD phenotypes, and also identify imaging features associated with previously identified genetic loci.

Abstract

Rationale: Chronic obstructive pulmonary disease (COPD) is defined by the presence of airflow limitation on spirometry, yet COPD subjects can have marked differences in CT imaging. These differences may be driven by genetic factors. We hypothesized that a genome-wide association study of quantitative imaging would identify loci not previously identified in analyses of COPD or spirometry. In addition, we sought to determine whether previously described genome-wide significant COPD and spirometric loci were associated with emphysema or airway phenotypes.

Objective: To identify genetic determinants of quantitative imaging phenotypes.

Methods: We performed a genome-wide association study on two quantitative emphysema and two quantitative airway imaging phenotypes in the COPDGene (non-Hispanic white and African-American), ECLIPSE, NETT, and GenKOLS studies; and on % gas trapping in COPDGene. We also examined specific loci reported as genome-wide significant for spirometric phenotypes related to airflow limitation or COPD.

Results: The total sample size across all cohorts was 12,031, of which 9,338 were from COPDGene. We identified five loci associated with emphysema-related phenotypes, one with airway-related phenotypes, and two with gas trapping. These loci included previously reported associations, including the *HHIP*, 15q25, and *AGER* loci, as well as novel associations near *SERPINA10* and *DLC1*. All previously reported COPD and a significant number of spirometric GWAS loci were at least nominally ($P < 0.05$) associated with either emphysema or airway phenotypes.

22 **Conclusions:** Genome-wide analysis may identify novel risk factors for quantitative imaging
23 characteristics in COPD, and also identify imaging features associated with previously identified
24 lung function loci.

Introduction

Chronic obstructive pulmonary disease (COPD) is a highly prevalent and morbid disease, defined by a simple measurement - the presence of irreversible airflow limitation on spirometry. Despite this simple clinical definition, COPD is a complex and heterogeneous disease with marked differences in the presence of key components that contribute to airflow obstruction in COPD – emphysema and airways disease (1). With the advent of standardized quantitative measurements, chest CT scans have become the prevalent method of characterizing lung parenchyma and airways in COPD(2).

Over the past several years, advances in image generation and analysis have led to studies demonstrating clinical and pathophysiologic relevance of these imaging measures. These include associations with spirometry(3, 4), respiratory symptoms(5), susceptibility to osteoporosis(6) and lung cancer(7), exacerbations(8), and lung function decline(9, 10).

The development of COPD is strongly influenced by genetic factors(11). Genetic variation is also an important determinant of emphysema and airway disease. Emphysema or airway imaging characteristics appear to be separately heritable(12, 13). Obstruction on pulmonary function can be seen in diseases predominantly involving the airway (in cystic fibrosis), or in those that involve the parenchyma through emphysema (alpha-1 antitrypsin deficiency and cutis laxa)(14). Previous genome-wide studies have identified variants associated with emphysema(15–17) or airway disease(18), though generally in smaller sample sizes or predominantly population-based subjects.

We hypothesized quantitative imaging reflects component disease processes leading to airflow obstruction in COPD, and could have genetic determinants not discovered by analyses

using lung function alone. To address this hypothesis, we performed a genome-wide association study of quantitative emphysema and airway phenotypes in current and former cigarette smokers with and without COPD. We additionally hypothesized genetic loci associated with spirometry related to airflow obstruction in general population samples or with COPD affection status would demonstrate an association with imaging phenotypes. Some of these results have been previously presented as an abstract(19).

Methods

Imaging measurements were available in COPDGene (NCT00608764. www.copd.org) non-Hispanic white and African-Americans, the Evaluation of COPD Longitudinally to Identify Predictive Surrogate Endpoints (ECLIPSE, SCO104960, NCT00292552, www.eclipse-copd.com), National Emphysema Treatment Trial (NETT), and GenKOLS (Genetics of COPD, Norway) study. Detailed descriptions including genotyping quality control, genotyping imputation, and quantitative imaging, have been previously published(5, 8, 20–27). All cohorts included only current or former smokers. COPDGene is a multicenter study including subjects of self-described non-Hispanic white or African-American ancestry and included subjects with and without COPD and with a range of spirometry. Subjects in the remaining studies were white. Controls had normal spirometry. Cases in the ECLIPSE and GenKOLS studies were at least GOLD spirometry grade 2 in severity. NETT cases had severe COPD ($FEV_1 < 45\%$ predicted) and were selected for the presence of emphysema.

Quantitative image analysis was performed on segmented CT chest images, using the number of voxels below -950 Hounsfield Units (%LAA-950) to estimate emphysema, and, alternatively, the Hounsfield Units at the 15th percentile of the density histogram (Perc15). The

airway wall area (Pi10) was the value for a hypothetical 10mm airway obtained by plotting a regression line of the square root of the airway wall area versus the airway internal perimeter(2). The wall area percent (WAP) was the percentage of the wall area compared to the total bronchial area for segmental and smaller airways (see Supplement). Percent gas trapping was measured at end-tidal exhalation and defined as the percent of lung voxels with < -856 HU(28).

We genotyped all subjects on Illumina platforms and imputed genotypes using MaCH and minimac(29) with 1000 Genomes Phase I v3 reference panels. We performed linear regression on each phenotype using residuals adjusted for age, sex, pack-years of smoking, current smoking status, and ancestry-based principal components. Imaging variables with marked non-normality were log-transformed (%LAA-950 and % gas trapping). COPDGene and ECLIPSE were additionally adjusted for CT scanner type. As airway measurements are not scaled to body size, we additionally adjusted for height. For gas trapping, a covariate for study center was also added to account for site-related technical variations in expiratory CT scans.

Results from all studies were combined into a meta-analysis. Given substantial heterogeneity within our studies, our primary analysis used a modified random-effects model(30). We also examined results using the standard fixed-effects model(31). As we hypothesized that emphysema and airway disease measured by quantitative CT may be causal for reduced lung function and COPD, our primary analyses included all subjects, with an additional analysis in cases only (including GOLD spirometry grade 1 for COPDGene subjects). To explore and control for the effect of ascertainment, we applied a method for analysis of secondary phenotype data within case-control association studies(32).

Additional methods are available in the Supplement.

Results

Genome-wide association of five quantitative imaging phenotypes

Baseline characteristics of subjects in each cohort are shown in **Table 1**. The total sample size across all cohorts was 12,031. Genome-wide significant results from the modified random-effects meta-analysis are shown in **Table 2**. Loci with prior evidence of association with COPD, lung function, and / or emphysema – *HHIP*, *CHRNA3/5/IREB2*, and *AGER* – were the most significant associations with %LAA-950. We also identified additional associations at genome-wide significance ($P < 5 \times 10^{-8}$) near *DLC1* and *SERPINA10*. An association near *CHRNA4* was just below genome-wide significance (rs183345681, $P = 1.8 \times 10^{-7}$). An analysis of Perc15 also identified the *DLC1* and *HHIP* loci associations.

In our analysis of airway phenotypes, no association reached genome-wide significance for Pi10. One result for wall area percent yielded $P < 5 \times 10^{-8}$ (rs142200419); however, this association was markedly attenuated in the fixed effects meta-analysis, due to effects in the opposite directions in one of the cohorts (Table S1). For the association analysis of gas trapping in COPDGene, the *AGER* and *LINC00310/KCNE2* loci achieved significance. No genome-wide significant results were identified in any of the case-only analyses (**Table S2**). For the regions yielding genome-wide significance in all subjects, we additionally examined results from an analysis accounting for ascertainment in COPDGene and GenKOLS, and including cases only from ECLIPSE (due to the small number of controls in this cohort). P-values obtained using this method(32) (**Table S1**) were generally only slightly less significant, with the possible exception of *HHIP* and *CHRNA3*, suggesting that overall our results were not simply driven by an association with case-control status. Results in cases and controls separately and, for loci not

previously described as genome-wide significant in COPD, a case-control analysis, are shown in
Tables S3 and S4.

The association with %LAA-950 near *SERPINA10* is also near *SERPINA1*, variants in which are the cause of alpha-1 antitrypsin deficiency. The most common form of severe alpha-1 antitrypsin deficiency is due to homozygosity for the Z allele, rs28929474. This variant was imputed with relatively high quality ($R_{sq} > 0.9$ in all white cohorts; 0.66 in COPDGene African-Americans). We examined the imputed rs28929474 in all cohorts, and did not find any ZZ subjects in NETT and GenKOLS; in COPDGene, seven non-Hispanic white ZZ subjects had been genotyped and subsequently excluded from analyses after *SERPINA1* genotyping (Foreman, In Preparation). All seven of these subjects were correctly identified with imputed genotypes. Linkage disequilibrium exists between our top associated SNP at this locus, rs45505795, and rs28929474 ($D' 0.7$, $r^2 = 0.295$). To determine if the association with rs45505795 could be accounted for by rs28929474, we performed a meta-analysis conditioned on rs28929474. The resulting P-value was 0.007, demonstrating that rs28929474 accounts for some, but not all, of the association signal. While known or identified ZZ homozygotes were excluded from COPDGene, NETT, and GenKOLS, ECLIPSE excluded only known alpha-1 deficient subjects. We identified six putative ZZ subjects in ECLIPSE. To determine whether the association signal in ECLIPSE was driven by the presence of these six subjects, we repeated the association analysis after dropping these subjects and found the P-value was slightly attenuated but remained significant ($P = 0.0018$), consistent overall with an increased risk of emphysema among MZ carriers.

To further explore the potential functional consequences of individual loci described in this study, we searched for evidence of functional impact using existing data sources. Of the loci

described in this study not previously associated with COPD, one was a cis-eQTL in lung – rs55706246 near *LINC00310* was in modest LD ($r^2 = 0.24$) with rs2834438, an eQTL for *KCNE2* ($p = 3.1 \times 10^{-7}$)(33). Using GWAS3D, the top-scoring variant at the *DLCI* locus was rs58863591, which had active enhancer marks (H3K4me1 and DNase hypersensitivity) and potential long-range interactions upstream of *DLCI* and near *SENP2*(34).

We also sought to determine whether the group of top (most significant) markers for each analysis ($P < 1 \times 10^{-6}$) could yield to insights about cell types based on regulatory data ENCODE(35). In the emphysema analysis, cell type enhancer enrichment from analysis of %LAA-950 among all subjects included enhancers in umbilical vein endothelial cells (Huvec, $P = 6.0 \times 10^{-4}$) and DNase I hypersensitivity sites in several types of endothelial cells ($P = 6.6 \times 10^{-3}$ to 0.03 for pulmonary artery endothelial cells (HPAEC) and adult blood, adult lymphatic, and neonatal lymphatic microvascular endothelial cells (HMVEC)). We found similar findings for the Perc15 analysis, with the strongest DNase enrichment for pulmonary artery endothelial cells ($P=0.017$). For the airway phenotypes, we found modest evidence for enrichment for enhancers K562 (leukemia) and HSMM (skeletal muscle) cell lines ($P = 0.02$) and DNase enrichment in CD14+ monocytes ($P = 0.04$).

We also sought to determine whether our results were consistent with a set of genes more likely to act within a specific gene sets or pathways. Top-ranked results identified several individual potential pathways of interest, including the toll-like receptor and phosphoinositide 3-kinase pathways (iGSEA4GWAS(36)) and telomere maintenance (INRICH(37)) for the %LAA-950 analyses. Gene sets that appeared to overlap between top-ranked sets among different methods included regulation of apoptosis, isoprenoid biosynthetic process, nicotinic

acetylcholine channel activity, actin cytoskeleton, and B-cell receptor signaling for emphysema GWAS; and for airway, WNT signaling and muscle contraction.

Associations at loci previously identified in association with COPD or COPD-related spirometric phenotypes

Genome-wide association studies have identified multiple variants associated with COPD(23–26, 38) or measures of lung function(39–41). We sought to determine whether there was evidence these variants might have an effect on quantitative imaging phenotypes, even if they did not reach genome-wide significance. After excluding loci previously associated in these cohorts with COPD, we found a strong enrichment in nominally significant (P -value < 0.05) loci among the two emphysema and two imaging phenotypes ($P = 4.9 \times 10^{-9}$), suggesting many of these variants may also affect quantitative imaging measurements. We further classified these variants into those showing a stronger association (by one-sided P -value) with emphysema- or airway-related phenotypes, assigning directionality such that the risk allele for COPD or reduced lung function demonstrated greater emphysema or increased airway wall thickness (**Table 3**). Enrichment for nominally significant P -values appeared to be greater among markers associated with quantitative emphysema ($P = 1.9 \times 10^{-6}$) versus those associated with airway wall thickness ($P=1.3 \times 10^{-3}$).

We next examined regulatory patterns using Haploreg(35) in variants classified as either emphysema or airway-associated identified in **Tables 2 & 3**. ‘Emphysema’ variants were modestly enriched for enhancers seen in hepatocellular carcinoma (HepG2, $P=0.05$), while those more strongly associated with airway phenotypes were enriched for enhancers from lung fibroblasts (NHLEF) and epidermal keratinocytes (NHEK, $P=0.03$ to 0.04). Both analyses were

enriched for mammary epithelial cells (HMEC, $P=2.5 \times 10^{-4}$ to 1.6×10^{-3}) and umbilical vein endothelial cells (Huvec, $P=0.02$ to 0.03). The most significant DNase enrichment for emphysema-associated variants was lung-derived lymphatic microvascular endothelial cells (HMVEC-LLy; $P 8 \times 10^{-4}$), while top results for airway-associated variants were embryonic lung fibroblasts (WI-38), mammary fibroblasts (HMF), and small airway epithelial cells (SAEC; $P 3.6-6.6 \times 10^{-4}$). Emphysema-associated DNase results were not significant in the airway results, and vice versa.

Discussion

In a genome-wide association study of quantitative imaging phenotypes in smokers with and without COPD, we identified genome-wide significant associations with loci previously shown to be associated with COPD or with spirometric measures related to airflow limitation, including the 15q25, *HHIP*, and *AGER* loci, the latter also identified in association with emphysema in a general population sample(15) and with emphysema and sRAGE levels in COPD(42). We also describe a genome-wide association with emphysema and variants near *SERPINA10*, and show that this association is in strong linkage disequilibrium with the Z-allele of *SERPINA1*, and not due the presence of PI ZZ individuals. This report is thus consistent with other reports showing an increased risk of airflow limitation for subjects with PI MZ(43, 44) and emphasizes the role of alpha-1 antitrypsin in the pathogenesis of COPD and emphysema in a broader group of patients.

One of our top associations with emphysema (both for %LAA-950 and Perc15) was a novel locus, located in the gene *DLC1* (deleted in liver cancer 1). *DLC1* frequently undergoes loss of heterozygosity or epigenetic silencing in solid cancers, including lung cancers(45). *DLC1*

appears to inhibit cell growth and increases apoptosis(46), and act as a tumor suppressor through the RhoGAP-dependent and RhoGAP-independent activity(47). *DLC1* is highly expressed in the lung(48, 49). In a study of regional emphysema, *DLC1* expression showed a trend towards decreased expression with an increase in the mean linear intercept(50) (nominal P-value, 0.04). Recently, a locus in *DLC1* was described in association with smoking behavior in African-Americans(51). We found a trend towards association with current smoking at this locus in COPDGene African-Americans ($P = 0.06-0.07$). However, we found no association with pack-years of smoking ($P > 0.49$). In addition, *DLC1* SNPs in this study are approximately 200kb away and not in linkage disequilibrium with our reported *DLC1* loci ($r^2 < 0.004$ in COPDGene African-Americans), and we found no consistent evidence of effect on either pack-years or current smoking at either locus in other cohorts. We also note an additional association near *CHRNA4* just below genome-wide significance. Previous studies have identified associations with smoking behavior in this region(52, 53), though previously described variants do not appear to be in strong LD with our identified variant. Additional studies will be needed to confirm our associations and determine their relationship to cigarette smoking.

We also examined variants previously identified at genome-wide significance in association with COPD or spirometric measures related to airflow obstruction. Most of these loci were at least nominally significantly ($P < 0.05$) associated with one or more quantitative CT phenotypes. Many appeared to have stronger associations with either quantitative emphysema or airway phenotypes. These findings suggest that genetic determinants of lung function in the general population may influence emphysema or airway disease, and are consistent with the hypothesis that there may be variants affecting airflow obstruction in different ways detectable by quantitative imaging.

In addition to examining individual loci, our study also explores the relevance of groups of markers that may not reach genome-wide significance. An analysis of gene sets provides supportive evidence for biological mechanisms previously been implicated in COPD, including telomere maintenance(54–57), phosphoinositide-3-kinase(58, 59), actin organization, and B-cell receptor signaling(50). An exploratory analysis of regulatory regions from ENCODE identified enrichment for endothelial cells. In animal models, targeted disruption of endothelial cells through genetic or immune mechanisms leading to apoptosis can lead to emphysema(60–62). Endothelial cell apoptosis has been seen in emphysematous human tissue(60) and endothelial microparticles, a marker for apoptosis, were related to emphysema in the MESA study(63). In contrast to prior work(16), we did not see an enrichment for fibroblasts from our quantitative emphysema analyses, but did see such enrichment in our airway-related lung function analysis.

Emphysema and airway disease are important components of COPD. We used automated and standardized measurements, available on a large number of subjects and free of inter-reader variation. We performed an analysis including all subjects in an effort to maximize power, and applied a method to account for ascertainment based on case-control status. However, due to the high correlation of disease status with imaging characteristics, we cannot rule out a degree of confounding for some of our associations. Although we performed five association analyses, we reported unadjusted P-values as our phenotypes are correlated, and some of our findings are seen in multiple phenotypes. Quantitative imaging can be affected by factors not related to intrinsic lung pathology, such as degree of inflation, obesity, smoking, and characteristics of individual CT scanners(5, 64, 65). Our decision to adjust for specific covariates was based on a desire to maximize findings of genetic analysis by controlling for the influence of age, smoking, and effects of individual scanners, yet allowing for genetic effects

that may affect disease processes contributing to more than one characteristic (e.g., low BMI and emphysema(66)). Ultimately, our findings will require replication, ideally in additional large cohorts that include a range of severity of COPD.

Our analysis also included studies with different imaging protocols, proportions of severity of disease, and racial groups. Thus, despite our large sample size, these factors may have resulted in a reduction in statistical power. We attempted to at least partially address this issue by using a method(30) that can improve power in the setting of heterogeneity. While most of the P-values from this method were very similar to those using standard fixed-effects models, this method resulted in *AGER* reaching genome-wide significance, consistent with prior studies. Our study is unable to address several causes of potential heterogeneity. Genetic factors may be specific to racial / ethnic groups(15). Technical factors may be less likely to influence reads by radiologists or semi-supervised methods and may explain why we were unable to replicate previous findings based on these approaches(16, 17). These factors, as well as differing proportions of severity of disease, may also indicate why we were unable to replicate findings from a recently reported analysis of airway wall thickness(18). Chest CT scans contain a wealth of data, and current measures of overall lung density or airway wall measurements do not adequately represent all relevant features. Efforts to expand and standardize radiologist interpretation and novel computational and machine learning-based methods may improve the ability to detect genetic effects.

Our work also demonstrates that previously described genetic associations with lung function in the population appear to influence airway or emphysema phenotypes. Using data from the ENCODE project, we identified non-overlapping enrichment of regulatory regions for our two sets of analyses. Our results are consistent with the hypothesis that emphysema and

airway imaging characteristics may be driven by different pathogenic processes and genetic factors(12). However, lung function, disease status, and imaging features are all correlated, and the relationship between specific imaging features is potentially complex(67). Our relative preponderance of associations with quantitative emphysema compared to airway, for example, may reflect the stronger correlation between lung function and our quantitative emphysema measurements or technical factors that affect airway measurements(67, 68). Our sets, particularly for ‘airway’ were loosely defined, and included results not reaching a nominal level of significance. Additional analytic methods, such as causal modeling, may help clarify the relationships between genetic variants, lung function, and CT imaging. Ultimately, however, the specific effects of individual variants will need to be determined by careful functional studies.

Differences in susceptibility to and phenotypic heterogeneity in COPD remain poorly understood. Despite their limitations, genome-wide association studies are currently the most powerful method to identify novel genetic risk factors for this complex and heterogeneous disease. Our analysis reflects a coordinated effort across multiple studies and to our knowledge is the largest genome-wide analysis of quantitative pulmonary imaging reported to date, and the first to include a substantial number of subjects with COPD. Our work identifies several genetic loci that may influence specific imaging phenotypes and identifies potential functional pathways and cell types through which these loci may exert their phenotypic effects. It also describes CT imaging phenotype-specific associations for loci previously implicated in GWAS for COPD or spirometric phenotypes related to COPD. Additional insights will result from increasing power; thus we anticipate a critical role for combining existing and upcoming studies using improved imaging phenotypes, to help unravel the complexity of pulmonary pathology in COPD.

References

1. Burrows B, Fletcher CM, Heard BE, Jones NL, Wootliff JS. The emphysematous and bronchial types of chronic airways obstruction. *Lancet* 1966;1:830–835.
2. Nakano Y, Muro S, Sakai H, Hirai T, Chin K, Tsukino M, Nishimura K, Itoh H, Paré PD, Hogg JC, Mishima M. Computed tomographic measurements of airway dimensions and emphysema in smokers. Correlation with lung function. *Am J Respir Crit Care Med* 2000;162:1102–8.
3. Mets OM, Buckens CFM, Zanen P, Isgum I, van Ginneken B, Prokop M, Gietema HA, Lammers J-WJ, Vliegenthart R, Oudkerk M, van Klaveren RJ, de Koning HJ, Mali WPTM, de Jong PA. Identification of chronic obstructive pulmonary disease in lung cancer screening computed tomographic scans. *JAMA* 2011;306:1775–81.
4. Washko GR, Criner GJ, Mohsenifar Z, Sciurba FC, Sharafkhaneh A, Make BJ, Hoffman EA, Reilly JJ. Computed tomographic-based quantification of emphysema and correlation to pulmonary function and mechanics. *COPD*, 2008/06/24 ed. 2008;5:177–186.
5. Grydeland TB, Dirksen A, Coxson HO, Eagan TML, Thorsen E, Pillai SG, Sharma S, Eide GE, Gulsvik A, Bakke PS. Quantitative computed tomography measures of emphysema and airway wall thickness are related to respiratory symptoms. *Am J Respir Crit Care Med* 2010;181:353–9.
6. Bon J, Fuhrman CR, Weissfeld JL, Duncan SR, Branch RA, Chang C-CH, Zhang Y, Leader JK, Gur D, Greenspan SL, Sciurba FC. Radiographic Emphysema Predicts Low Bone Mineral Density in a Tobacco-exposed Cohort. *Am J Respir Crit Care Med* 2011;183:885–90.
7. Wilson DO, Weissfeld JL, Balkan A, Schragin JG, Fuhrman CR, Fisher SN, Wilson J, Leader JK, Siegfried JM, Shapiro SD, Sciurba FC. Association of radiographic emphysema and airflow obstruction with lung cancer. *Am J Respir Crit Care Med*, 2008/06/21 ed. 2008;178:738–744.
8. Han MK, Kazerooni EA, Lynch DA, Liu LX, Murray S, Curtis JL, Criner GJ, Kim V, Bowler RP, Hanania NA, Anzueto AR, Make BJ, Hokanson JE, Crapo JD, Silverman EK, Martinez FJ, Washko GR, The COPDGene Investigators F. Chronic obstructive pulmonary disease exacerbations in the COPDGene study: associated radiologic phenotypes. *Radiology* 2011;261:274–82.
9. Mohamed Hoesein FAA, de Hoop B, Zanen P, Gietema H, Kruitwagen CLJJ, van Ginneken B, Isgum I, Mol C, van Klaveren RJ, Dijkstra AE, Groen HJM, Boezen HM, Postma DS, Prokop M, Lammers J-WJ. CT-quantified emphysema in male heavy smokers: association with lung function decline. *Thorax* 2011;66:782–7.

- 329 10. Nishimura M, Makita H, Nagai K, Konno S, Nasuhara Y, Hasegawa M, Shimizu K,
330 Betsuyaku T, Ito YM, Fuke S, Igarashi T, Akiyama Y, Ogura S. Annual change in
331 pulmonary function and clinical phenotype in chronic obstructive pulmonary disease. *Am*
332 *J Respir Crit Care Med* 2012;185:44–52.
- 333 11. Ingebrigtsen T, Thomsen SF, Vestbo J, van der Sluis S, Kyvik KO, Silverman EK,
334 Svartengren M, Backer V. Genetic influences on Chronic Obstructive Pulmonary Disease
335 - a twin study. *Respir Med* 2010;104:1890–5.
- 336 12. Patel BD, Coxson HO, Pillai SG, Agustí AGN, Calverley PMA, Donner CF, Make BJ,
337 Müller NL, Rennard SI, Vestbo J, Wouters EFM, Hiorns MP, Nakano Y, Camp PG,
338 Nasute Fauerbach P V, Screaton NJ, Campbell EJ, Anderson WH, Paré PD, Levy RD,
339 Lake SL, Silverman EK, Lomas DA. Airway wall thickening and emphysema show
340 independent familial aggregation in chronic obstructive pulmonary disease. *Am J Respir*
341 *Crit Care Med* 2008;178:500–5.
- 342 13. Zhou JJ, Cho MH, Castaldi PJ, Hersh CP, Silverman EK, Laird NM. Heritability of COPD
343 and Related Phenotypes in Smokers. *Am J Respir Crit Care Med*
344 2013;doi:10.1164/rccm.201302-0263OC.
- 345 14. Urban Z, Gao J, Pope FM, Davis EC. Autosomal dominant cutis laxa with severe lung
346 disease: synthesis and matrix deposition of mutant tropoelastin. 2005;124:1193–1199.
- 347 15. Manichaikul A, Hoffman EA, Smolonska J, Gao W, Cho MH, Baumhauer H, Budoff M,
348 Austin JHM, Washko GR, Carr JJ, Kaufman JD, Pottinger T, Powell CA, Wijmenga C,
349 Zanen P, Groen HJM, Postma DS, Wanner A, Rouhani FN, Brantly ML, Powell R, Smith
350 BM, Rabinowitz D, Raffel LJ, Hinckley Stukovsky KD, Crapo JD, Beaty TH, Hokanson
351 JE, Silverman EK, *et al.* Genome-wide study of percent emphysema on computed
352 tomography in the general population. The Multi-Ethnic Study of Atherosclerosis
353 Lung/SNP Health Association Resource Study. *Am J Respir Crit Care Med*
354 2014;189:408–18.
- 355 16. Castaldi PJ, Cho MH, Estépar RSJ, McDonald M-LN, Laird N, Beaty TH, Washko G,
356 Crapo JD, Silverman EK. Genome-Wide Association Identifies Regulatory Loci
357 Associated with Distinct Local Histogram Emphysema Patterns. *Am J Respir Crit Care*
358 *Med* 2014;doi:10.1164/rccm.201403-0569OC.
- 359 17. Kong X, Cho MH, Anderson W, Coxson HO, Muller N, Washko G, Hoffman EA, Bakke
360 P, Gulsvik A, Lomas DA, Silverman EK, Pillai SG. Genome-wide association study
361 identifies BICD1 as a susceptibility gene for emphysema. *Am J Respir Crit Care Med*
362 2011;183:43–9.
- 363 18. Dijkstra AE, Postma DS, van Ginneken B, Wielpütz MO, Schmidt M, Becker N,
364 Owsijewitsch M, Kauczor H-U, de Koning HJ, Lammers JW, Oudkerk M, Brandsma C-A,
365 Bossé Y, Nickle DC, Sin DD, Hiemstra PS, Wijmenga C, Smolonska J, Zanen P, Vonk
366 JM, van den Berge M, Boezen HM, Groen HJM. Novel Genes for Airway Wall Thickness

- 367 Identified with Combined Genome Wide Association and Expression Analyses. *Am J*
 368 *Respir Crit Care Med* 2014;doi:10.1164/rccm.201405-0840OC.
- 369 19. Cho MH, Castaldi P, Hersh CP, Barr RG, Tal-Singer R, Bakke P, Gulsvik A, Van Beek
 370 EJ, Coxson HO, Lynch DA, Washko GR, Laird N, Crapo J, Beaty T, Silverman EK. A
 371 Genome-Wide Association Study of Emphysema and Airway Quantitative Imaging
 372 Phenotypes. *Am Thorac Soc Int Conf* 2015. p. A2849.
- 373 20. Regan EA, Hokanson JE, Murphy JR, Make B, Lynch DA, Beaty TH, Curran-Everett D,
 374 Silverman EK, Crapo JD. Genetic epidemiology of COPD (COPDGene) study design.
 375 *COPD*, 2010/03/11 ed. 7:32–43.
- 376 21. Vestbo J, Anderson W, Coxson HO, Crim C, Dawber F, Edwards L, Hagan G, Knobil K,
 377 Lomas DA, MacNee W, Silverman EK, Tal-Singer R. Evaluation of COPD
 378 Longitudinally to Identify Predictive Surrogate End-points (ECLIPSE). *Eur Respir J*
 379 2008;31:869–73.
- 380 22. Fishman A, Martinez F, Naunheim K, Piantadosi S, Wise R, Ries A, Weinmann G, Wood
 381 DE. A randomized trial comparing lung-volume-reduction surgery with medical therapy
 382 for severe emphysema. *N Engl J Med* 2003;348:2059–73.
- 383 23. Pillai SG, Ge D, Zhu G, Kong X, Shianna K V, Need AC, Feng S, Hersh CP, Bakke P,
 384 Gulsvik A, Ruppert A, Lodrup Carlsen KC, Roses A, Anderson W, Rennard SI, Lomas
 385 DA, Silverman EK, Goldstein DB. A genome-wide association study in chronic
 386 obstructive pulmonary disease (COPD): identification of two major susceptibility loci.
 387 *PLoS Genet*, 2009/03/21 ed. 2009;5:e1000421.
- 388 24. Cho MH, Boutaoui N, Klanderman BJ, Sylvia JS, Ziniti JP, Hersh CP, DeMeo DL,
 389 Hunninghake GM, Litonjua AA, Sparrow D, Lange C, Won S, Murphy JR, Beaty TH,
 390 Regan EA, Make BJ, Hokanson JE, Crapo JD, Kong X, Anderson WH, Tal-Singer R,
 391 Lomas DA, Bakke P, Gulsvik A, Pillai SG, Silverman EK. Variants in FAM13A are
 392 associated with chronic obstructive pulmonary disease. *Nat Genet*, 2010/02/23 ed.
 393 2010;42:200–202.
- 394 25. Cho MH, Castaldi PJ, Wan ES, Siedlinski M, Hersh CP, Demeo DL, Himes BE, Sylvia
 395 JS, Klanderman BJ, Ziniti JP, Lange C, Litonjua AA, Sparrow D, Regan EA, Make BJ,
 396 Hokanson JE, Murray T, Hetmanski JB, Pillai SG, Kong X, Anderson WH, Tal-Singer R,
 397 Lomas DA, Coxson HO, Edwards LD, MacNee W, Vestbo J, Yates JC, Agusti A, *et al.* A
 398 genome-wide association study of COPD identifies a susceptibility locus on chromosome
 399 19q13. *Hum Mol Genet* 2012;21:947–57.
- 400 26. Cho MH, McDonald M-LN, Zhou X, Mattheisen M, Castaldi PJ, Hersh CP, Demeo DL,
 401 Sylvia JS, Ziniti J, Laird NM, Lange C, Litonjua AA, Sparrow D, Casaburi R, Barr RG,
 402 Regan EA, Make BJ, Hokanson JE, Lutz S, Dudenkov TM, Farzadegan H, Hetmanski JB,
 403 Tal-Singer R, Lomas DA, Bakke P, Gulsvik A, Crapo JD, Silverman EK, Beaty TH. Risk

404 loci for chronic obstructive pulmonary disease: a genome-wide association study and
405 meta-analysis. *lancet Respir Med* 2014;2:214–25.

406 27. DeMeo DL, Hersh CP, Hoffman EA, Litonjua AA, Lazarus R, Sparrow D, Benditt JO,
407 Criner G, Make B, Martinez FJ, Scanlon PD, Sciurba FC, Utz JP, Reilly JJ, Silverman
408 EK. Genetic determinants of emphysema distribution in the national emphysema
409 treatment trial. *Am J Respir Crit Care Med* 2007;176:42–48.

410 28. Schroeder JD, McKenzie AS, Zach JA, Wilson CG, Curran-Everett D, Stinson DS,
411 Newell JD, Lynch DA. Relationships between airflow obstruction and quantitative CT
412 measurements of emphysema, air trapping, and airways in subjects with and without
413 chronic obstructive pulmonary disease. *AJR Am J Roentgenol* 2013;201:W460–70.

414 29. Howie B, Fuchsberger C, Stephens M, Marchini J, Abecasis GR. Fast and accurate
415 genotype imputation in genome-wide association studies through pre-phasing. *Nat Genet*
416 2012;44:955–9.

417 30. Han B, Eskin E. Random-Effects Model Aimed at Discovering Associations in Meta-
418 Analysis of Genome-wide Association Studies. *Am J Hum Genet* 2011;88:586–98.

419 31. De Bakker PIW, Ferreira M a R, Jia X, Neale BM, Raychaudhuri S, Voight BF. Practical
420 aspects of imputation-driven meta-analysis of genome-wide association studies. *Hum Mol*
421 *Genet* 2008;17:R122–8.

422 32. Lin DY, Zeng D. Proper analysis of secondary phenotype data in case-control association
423 studies. *Genet Epidemiol* 2009;33:256–65.

424 33. Hao K, Bossé Y, Nickle DC, Paré PD, Postma DS, Laviolette M, Sandford A, Hackett TL,
425 Daley D, Hogg JC, Elliott WM, Couture C, Lamontagne M, Brandsma C-A, van den
426 Berge M, Koppelman G, Reicin AS, Nicholson DW, Malkov V, Derry JM, Suver C, Tsou
427 JA, Kulkarni A, Zhang C, Vessey R, Opiteck GJ, Curtis SP, Timens W, Sin DD. Lung
428 eQTLs to help reveal the molecular underpinnings of asthma. *PLoS Genet*
429 2012;8:e1003029.

430 34. Li MJ, Wang LY, Xia Z, Sham PC, Wang J. GWAS3D: Detecting human regulatory
431 variants by integrative analysis of genome-wide associations, chromosome interactions
432 and histone modifications. *Nucleic Acids Res* 2013;41:W150–8.

433 35. Ward LD, Kellis M. HaploReg: a resource for exploring chromatin states, conservation,
434 and regulatory motif alterations within sets of genetically linked variants. *Nucleic Acids*
435 *Res* 2012;40:D930–4.

436 36. Zhang K, Cui S, Chang S, Zhang L, Wang J. i-GSEA4GWAS: a web server for
437 identification of pathways/gene sets associated with traits by applying an improved gene
438 set enrichment analysis to genome-wide association study. *Nucleic Acids Res*, 2010/05/04
439 ed. 2010;38:W90–5.

- 440 37. Lee PH, O'Dushlaine C, Thomas B, Purcell SM. INRICH: interval-based enrichment
441 analysis for genome-wide association studies. *Bioinformatics* 2012;28:1797–9.
- 442 38. Wilk JB, Shrine NRG, Loehr LR, Zhao JH, Manichaikul A, Lopez LM, Smith AV,
443 Heckbert SR, Smolonska J, Tang W, Loth DW, Curjuric I, Hui J, Cho MH, Latourelle JC,
444 Henry AP, Aldrich M, Bakke P, Beaty TH, Bentley AR, Borecki IB, Brusselle GG,
445 Burkart KM, Chen T, Couper D, Crapo JD, Davies G, Dupuis J, Franceschini N, *et al.*
446 Genome-wide association studies identify CHRNA5/3 and HTR4 in the development of
447 airflow obstruction. *Am J Respir Crit Care Med* 2012;186:622–32.
- 448 39. Soler Artigas M, Loth DW, Wain L V, Gharib SA, Obeidat M, Tang W, Zhai G, Zhao JH,
449 Smith AV, Huffman JE, Albrecht E, Jackson CM, Evans DM, Cadby G, Fornage M,
450 Manichaikul A, Lopez LM, Johnson T, Aldrich MC, Aspelund T, Barroso I, Campbell H,
451 Cassano PA, Couper DJ, Eiriksdottir G, Franceschini N, Garcia M, Gieger C, Gislason
452 GK, *et al.* Genome-wide association and large-scale follow up identifies 16 new loci
453 influencing lung function. *Nat Genet* 2011;43:1082–90.
- 454 40. Hancock DB, Artigas MS, Gharib SA, Henry A, Manichaikul A, Ramasamy A, Loth DW,
455 Imboden M, Koch B, McArdle WL, Smith A V, Smolonska J, Sood A, Tang W, Wilk JB,
456 Zhai G, Zhao JH, Aschard H, Burkart KM, Curjuric I, Eijgelsheim M, Elliott P, Gu X,
457 Harris TB, Janson C, Homuth G, Hysi PG, Liu JZ, Loehr LR, *et al.* Genome-wide joint
458 meta-analysis of SNP and SNP-by-smoking interaction identifies novel loci for pulmonary
459 function. *PLoS Genet* 2012;8:e1003098.
- 460 41. Repapi E, Sayers I, Wain L V, Burton PR, Johnson T, Obeidat M, Zhao JH, Ramasamy A,
461 Zhai G, Vitart V, Huffman JE, Igl W, Albrecht E, Deloukas P, Henderson J, Granell R,
462 McArdle WL, Rudnicka AR, Barroso I, Loos RJ, Wareham NJ, Mustelin L, Rantanen T,
463 Surakka I, Imboden M, Wichmann HE, Grkovic I, Jankovic S, Zgaga L, *et al.* Genome-
464 wide association study identifies five loci associated with lung function. *Nat Genet*,
465 2009/12/17 ed. 2009;42:36–44.
- 466 42. Cheng DT, Kim DK, Cockayne DA, Belousov A, Bitter H, Cho MH, Duvoix A, Edwards
467 LD, Lomas DA, Miller BE, Reynaert N, Tal-Singer R, Wouters EFM, Agustí A, Fabbri
468 LM, Rames A, Visvanathan S, Rennard SI, Jones P, Parmar H, MacNee W, Wolff G,
469 Silverman EK, Mayer RJ, Pillai SG. Systemic soluble receptor for advanced glycation
470 endproducts is a biomarker of emphysema and associated with AGER genetic variants in
471 patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*
472 2013;188:948–57.
- 473 43. Molloy K, Hersh CP, Morris VB, Carroll TP, O'Connor CA, Lasky-Su JA, Greene CM,
474 O'Neill SJ, Silverman EK, McElvaney NG. Clarification of the risk of chronic obstructive
475 pulmonary disease in α 1-antitrypsin deficiency PiMZ heterozygotes. *Am J Respir Crit*
476 *Care Med* 2014;189:419–27.
- 477 44. Sørheim I-C, Bakke P, Gulsvik A, Pillai SG, Johannessen A, Gaarder PI, Campbell EJ,
478 Agustí A, Calverley PMA, Donner CF, Make BJ, Rennard SI, Vestbo J, Wouters EFM,

- 479 Paré PD, Levy RD, Coxson HO, Lomas DA, Hersh CP, Silverman EK. α_1 -Antitrypsin
480 protease inhibitor MZ heterozygosity is associated with airflow obstruction in two large
481 cohorts. *Chest* 2010;138:1125–32.
- 482 45. Liao Y-C, Lo SH. Deleted in liver cancer-1 (DLC-1): a tumor suppressor not just for liver.
483 *Int J Biochem Cell Biol* 2008;40:843–7.
- 484 46. Zhou X, Thorgeirsson SS, Popescu NC. Restoration of DLC-1 gene expression induces
485 apoptosis and inhibits both cell growth and tumorigenicity in human hepatocellular
486 carcinoma cells. *Oncogene* 2004;23:1308–13.
- 487 47. Healy KD, Hodgson L, Kim T-Y, Shutes A, Maddileti S, Juliano RL, Hahn KM, Harden
488 TK, Bang Y-J, Der CJ. DLC-1 suppresses non-small cell lung cancer growth and invasion
489 by RhoGAP-dependent and independent mechanisms. *Mol Carcinog* 2008;47:326–37.
- 490 48. Uhlen M, Oksvold P, Fagerberg L, Lundberg E, Jonasson K, Forsberg M, Zwahlen M,
491 Kampf C, Wester K, Hober S, Wernerus H, Björling L, Ponten F. Towards a knowledge-
492 based Human Protein Atlas. *Nat Biotechnol* 2010;28:1248–50.
- 493 49. Lonsdale J, Thomas J, Salvatore M, Phillips R, Lo E, Shad S, Hasz R, Walters G, Garcia
494 F, Young N, Foster B, Moser M, Karasik E, Gillard B, Ramsey K, Sullivan S, Bridge J,
495 Magazine H, Syron J, Fleming J, Siminoff L, Traino H, Mosavel M, Barker L, Jewell S,
496 Rohrer D, Maxim D, Filkins D, Harbach P, *et al.* The Genotype-Tissue Expression
497 (GTEx) project. *Nat Genet*, 2013/05/30 ed. 2013;45:580–585.
- 498 50. Campbell JD, McDonough JE, Zeskind JE, Hackett TL, Pechkovsky D V, Brandsma C-A,
499 Suzuki M, Gosselink J V, Liu G, Alekseyev YO, Xiao J, Zhang X, Hayashi S, Cooper JD,
500 Timens W, Postma DS, Knight DA, Lenburg ME, Hogg JC, Spira A. A gene expression
501 signature of emphysema-related lung destruction and its reversal by the tripeptide GHK.
502 *Genome Med* 2012;4:67.
- 503 51. Gelernter J, Kranzler HR, Sherva R, Almasy L, Herman AI, Koesterer R, Zhao H, Farrer
504 LA. Genome-wide association study of nicotine dependence in American populations:
505 identification of novel risk loci in both African-Americans and European-Americans. *Biol*
506 *Psychiatry* 2015;77:493–503.
- 507 52. Lazary J, Dome P, Csala I, Kovacs G, Faludi G, Kaunisto M, Dome B. Massive
508 withdrawal symptoms and affective vulnerability are associated with variants of the
509 CHRNA4 gene in a subgroup of smokers. In: Chen L, editor. *PLoS One* 2014;9:e87141.
- 510 53. Han S, Yang B-Z, Kranzler HR, Oslin D, Anton R, Gelernter J. Association of CHRNA4
511 polymorphisms with smoking behavior in two populations. *Am J Med Genet B*
512 *Neuropsychiatr Genet* 2011;156B:421–9.
- 513 54. Savale L, Chaouat A, Bastuji-Garin S, Marcos E, Boyer L, Maitre B, Sarni M, Housset B,
514 Weitzenblum E, Matrat M, Le Corvoisier P, Rideau D, Boczkowski J, Dubois-Randé J-L,

- 515 Chouaid C, Adnot S. Shortened telomeres in circulating leukocytes of patients with
516 chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2009;179:566–71.
- 517 55. Tsuji T, Aoshiba K, Nagai A. Alveolar cell senescence in patients with pulmonary
518 emphysema. *Am J Respir Crit Care Med* 2006;174:886–93.
- 519 56. Alder JK, Guo N, Kembou F, Parry EM, Anderson CJ, Gorgy AI, Walsh MF, Sussan T,
520 Biswal S, Mitzner W, Tudor RM, Armanios M. Telomere Length is a Determinant of
521 Emphysema Susceptibility. *Am J Respir Crit Care Med* 2011;doi:10.1164/rccm.201103-
522 0520OC.
- 523 57. Nunes H, Monnet I, Kannengiesser C, Uzunhan Y, Valeyre D, Kambouchner M,
524 Naccache J-M. Is telomeropathy the explanation for combined pulmonary fibrosis and
525 emphysema syndrome?: report of a family with TERT mutation. *Am J Respir Crit Care*
526 *Med* 2014;189:753–4.
- 527 58. Barnes PJ. Emerging pharmacotherapies for COPD. *Chest*, 2008/12/09 ed.
528 2008;134:1278–1286.
- 529 59. Fang X, Li K, Tao X, Chen C, Wang X, Wang L, Wang DC, Zhang Y, Bai C, Wang X.
530 Effects of phosphoinositide 3-kinase on protease-induced acute and chronic lung
531 inflammation, remodeling, and emphysema in rats. *Chest* 2013;143:1025–35.
- 532 60. Kasahara Y, Tudor RM, Taraseviciene-Stewart L, Le Cras TD, Abman S, Hirth PK,
533 Waltenberger J, Voelkel NF. Inhibition of VEGF receptors causes lung cell apoptosis and
534 emphysema. 2000;106:1311–1319.
- 535 61. Taraseviciene-Stewart L, Scerbavicius R, Choe K-H, Moore M, Sullivan A, Nicolls MR,
536 Fontenot AP, Tudor RM, Voelkel NF. An animal model of autoimmune emphysema. *Am J*
537 *Respir Crit Care Med* 2005;171:734–42.
- 538 62. Giordano RJ, Lahdenranta J, Zhen L, Chukwueke U, Petrache I, Langley RR, Fidler IJ,
539 Pasqualini R, Tudor RM, Arap W. Targeted induction of lung endothelial cell apoptosis
540 causes emphysema-like changes in the mouse. *J Biol Chem* 2008;283:29447–60.
- 541 63. Thomashow MA, Shimbo D, Parikh MA, Hoffman EA, Vogel-Claussen J, Hueper K, Fu
542 J, Liu C-Y, Bluemke DA, Ventetuolo CE, Doyle MF, Barr RG. Endothelial microparticles
543 in mild chronic obstructive pulmonary disease and emphysema. The Multi-Ethnic Study
544 of Atherosclerosis Chronic Obstructive Pulmonary Disease study. *Am J Respir Crit Care*
545 *Med* 2013;188:60–8.
- 546 64. Stratelis G, Fransson SG, Schmekel B, Jakobsson P, Molstad S. High prevalence of
547 emphysema and its association with BMI: A study of smokers with normal spirometry.
548 2008/10/11 ed. 2008;1–7.doi:903492403 [pii] 10.1080/02813430802452732.

549 65. Hoffman EA, Ahmed FS, Baumhauer H, Budoff M, Carr JJ, Kronmal R, Reddy S, Barr
550 RG. Variation in the percent of emphysema-like lung in a healthy, nonsmoking
551 multiethnic sample. The MESA lung study. *Ann Am Thorac Soc* 2014;11:898–907.

552 66. Wan ES, Cho MH, Boutaoui N, Klanderman BJ, Sylvia JS, Ziniti JP, Won S, Lange C,
553 Pillai SG, Anderson WH, Kong X, Lomas DA, Bakke PS, Gulsvik A, Regan EA, Murphy
554 JR, Make BJ, Crapo JD, Wouters EF, Celli BR, Silverman EK, Demeo DL. Genome-Wide
555 Association Analysis of Body Mass in Chronic Obstructive Pulmonary Disease. *Am J*
556 *Respir Cell Mol Biol* 2010;doi:10.1165/rcmb.2010-0294OC.

557 67. Diaz AA, Come CE, Ross JC, San José Estépar R, Han MK, Loring SH, Silverman EK,
558 Washko GR. Association between airway caliber changes with lung inflation and
559 emphysema assessed by volumetric CT scan in subjects with COPD. *Chest* 2012;141:736–
560 44.

561 68. Smith BM, Hoffman EA, Rabinowitz D, Bleecker E, Christenson S, Couper D, Donohue
562 KM, Han MK, Hansel NN, Kanner RE, Kleerup E, Rennard S, Barr RG. Comparison of
563 spatially matched airways reveals thinner airway walls in COPD. The Multi-Ethnic Study
564 of Atherosclerosis (MESA) COPD Study and the Subpopulations and Intermediate
565 Outcomes in COPD Study (SPIROMICS). *Thorax* 2014;doi:10.1136/thoraxjnl-2014-
566 205160.

567 69. Repapi E, Sayers I, Wain L V, Burton PR, Johnson T, Obeidat M, Zhao JH, Ramasamy A,
568 Zhai G, Vitart V, Huffman JE, Igl W, Albrecht E, Deloukas P, Henderson J, Granell R,
569 McArdle WL, Rudnicka AR, Barroso I, Loos RJF, Wareham NJ, Mustelin L, Rantanen T,
570 Surakka I, Imboden M, Wichmann HE, Grkovic I, Jankovic S, Zgaga L, *et al.* Genome-
571 wide association study identifies five loci associated with lung function. *Nat Genet*
572 2010;42:36–44.

573 70. Wilk JB, Chen TH, Gottlieb DJ, Walter RE, Nagle MW, Brandler BJ, Myers RH, Borecki
574 IB, Silverman EK, Weiss ST, O'Connor GT. A genome-wide association study of
575 pulmonary function measures in the Framingham Heart Study. *Am J Respir Crit Care*
576 *Med*, 2009/03/21 ed. 2009;5:e1000429.

577 71. Hancock DB, Eijgelsheim M, Wilk JB, Gharib SA, Loehr LR, Marcianti KD,
578 Franceschini N, van Durme YMTA, Chen T-HH, Barr RG, Schabath MB, Couper DJ,
579 Brusselle GG, Psaty BM, van Duijn CM, Rotter JI, Uitterlinden AG, Hofman A, Punjabi
580 NM, Rivadeneira F, Morrison AC, Enright PL, North KE, Heckbert SR, Lumley T,
581 Stricker BHC, O'Connor GT, London SJ. Meta-analyses of genome-wide association
582 studies identify multiple loci associated with pulmonary function. *Nat Genet*, 2009/12/17
583 ed. 2009;42:45–52.

Tables

Table 1: Baseline characteristics of subjects with quantitative imaging phenotypes. Cases = GOLD Grade 2 or more severe (e.g. NETT) cases; Controls = GOLD 0 smoking controls; Non-cases: includes GOLD 0, 1, and PRISm subjects.

	COPDGene non-Hispanic Whites		COPDGene African Americans		ECLIPSE		NETT	GenKOLS (Norway)	
	Non-cases	Cases	Non-cases	Cases	Controls	Cases	Cases	Controls	Cases
n	3062	3243	2132	901	145	1393	332	406	417
Age	59.7 (8.6)	64.4 (8.3)	53 (6)	58.6 (8.1)	57.3 (9.4)	63.4 (7)	67.4 (5.9)	55.6 (9.4)	64.2 (9.3)
Pack-years	39.7 (21.5)	54.4 (27.5)	36.6 (20.5)	42 (23.1)	31.8 (26.6)	49.8 (26.7)	65.8 (30.8)	19.8 (14.1)	31 (18.2)
Sex (%Male)	1462 (47.7%)	1832 (56.5%)	1209 (56.7%)	497 (55.2%)	85 (58.6%)	911 (65.4%)	212 (63.9%)	216 (53.2%)	263 (63.1%)
Current smokers	1263 (41.2%)	1199 (37%)	1838 (86.2%)	595 (66%)	58 (40%)	480 (34.5%)	0	164 (40.4%)	210 (50.4%)
FEV₁, % predicted	91.3 (14.8)	57.4 (23)	92.2 (16.5)	59.5 (22)	108.6 (13.4)	47.4 (15.5)	28.2 (7.3)	94.9 (9.2)	52.5 (16.9)
%LAA-950	1.2 (0-26.9)	7.5 (0-61.9)	0.7 (0-35.8)	4.6 (0-61.2)	2.3 (0.1-14.2)	16.3 (0.1-58.7)	15 (0.3-49.9)	0.5 (0-34.4)	7 (0-53.2)
Perc15, HU	-909.9 (22.8)	-938.1 (26.8)	-893.4 (28.1)	-926.5 (32)	-906.2 (25.9)	-950.9 (25.9)	-949.7 (17.8)	-891.6 (26.3)	-932.8 (30.2)
Pi10, mm	3.64 (0.11)	3.69 (0.14)	3.69 (0.13)	3.73 (0.15)	4.34 (0.15)	4.41 (0.20)	4.58 (0.49)	4.76 (0.29)	4.94 (0.34)
Wall area percent (WAP)	60.2 (2.8)	62.3 (3.1)	61.2 (3.3)	62.9 (3.3)	63.2 (3.7)	65.6 (4.1)	73.2 (3.8)	74.8 (2.9)	76.1 (3)
Gas trapping, %	9.3 (0-83.4)	34 (0.1-87.8)	7.2 (0-70.5)	29.3 (0.2-85.2)					

Table 2: Genome-wide significant associations. %LAA-950: percent of low attenuation area less than -950 Hounsfield units; Perc15 - Hounsfield Units at the 15th percentile of the density histogram; WAP percentage of the wall area compared to the total bronchial area.

Phenotype	Chr	Marker Name	Closest Gene	Effect Allele	Allele Frequency		Modified Random Effects			Fixed Effects		
					Nhw	Aa	P value	Beta	SE	P value	Beta	Se
<i>Emphysema</i>												
%LAA-950	4	rs13141641	HHIP	T	0.59	0.89	1.7×10^{-12}	0.12	0.023	8.4×10^{-13}	0.12	0.018
	15	rs55676755	CHRNA3	C	0.63	0.84	2.4×10^{-9}	-0.11	0.017	1.4×10^{-9}	-0.11	0.017
	6	rs2070600	AGER	T	0.04	0.01	4.6×10^{-9}	-0.14	0.11	6.5×10^{-8}	-0.24	0.044
	8	rs75200691	DLC1	T	0.88	0.92	9.7×10^{-9}	0.15	0.026	5.7×10^{-9}	0.15	0.026
	14	rs45505795	SERPINA10	C	0.04	0.008	1.4×10^{-8}	-0.31	0.08	9.8×10^{-9}	-0.31	0.053
Perc 15	8	rs74834049	DLC1	A	0.12	0.08	6.0×10^{-10}	-3.4	0.54	3.3×10^{-10}	-3.4	0.54
	4	rs13141641	HHIP	T	0.59	0.89	8.4×10^{-10}	-2.2	0.39	4.7×10^{-10}	-2.2	0.36
Airway												
WAP	4	rs142200419	MIR2054	T	0.98	N/A	4.6×10^{-9}	0.24	1	8.8×10^{-5}	0.9	0.23
Gas trapping												
%	6	rs2070600	AGER	T	0.04	0.01	3.5×10^{-9}	-0.23	0.039	2.4×10^{-9}	-0.23	0.039
	21	rs55706246	LINC00310	A	0.11	0.03	1.3×10^{-8}	0.28	0.18	2.1×10^{-7}	0.15	0.029

Table 3: P-values for genetic variants previously reported in genome-wide association analyses(23–26, 39, 40, 69–71). The risk allele for spirometric phenotypes denotes the allele associated with a lower FEV₁ or FEV₁/FVC ratio, and thus would be expected to increase risk for COPD. The sign associated with the P-values denotes whether the direction of association is consistent with the direction for COPD (increase in %LAA-950, Pi10, wall area percent, or gas trapping; decrease in Perc15). In Table 3b, results are grouped by whether the smaller directional P-value was found in emphysema phenotypes (top) or airway-related phenotypes (bottom). Genome-wide significant loci from **Table 2** (e.g. *HHIP*) are not included here. All refers to all subjects, case refers to all cases (GOLD 1-4 or 2-4).

Table 3a: Variants from GWAS of moderate-to-severe or severe COPD

SNP	Chr	Locus	Risk Allele	Emphysema				Airway				Gas Trapping	
				%LAA-950		Perc15		Pi10		Wall Area Percent			
				All	Case	All	Case	All	Case	All	Case	All	Case
rs626750	11	<i>MMP12</i>	G	2x10 ⁻⁵	4x10⁻⁷	6x10 ⁻⁶	7x10 ⁻⁷	-0.1	-0	0.2	-0.1	0.008	0.1
rs4846480	1	<i>TGFB2</i>	A	2x10⁻⁶	3x10 ⁻⁵	1x10 ⁻⁴	5x10 ⁻⁴	-0.7	-0.4	0.2	-0.9	3x10 ⁻⁴	0.009
rs7937	19	<i>RAB4B</i>	T	2x10⁻⁶	0.03	6x10 ⁻⁵	0.03	0.9	-0.08	0.4	-0.04	9x10 ⁻⁴	0.2
rs754388	14	<i>RIN3</i>	C	3x10⁻⁵	0.1	5x10 ⁻⁵	0.04	0.4	-0.5	0.04	-0.6	0.003	0.1
rs7671167	4	<i>FAM13A</i>	T	3x10 ⁻⁴	0.3	2x10⁻⁴	0.07	0.6	-0.8	0.1	-0.5	9x10 ⁻⁵	0.6

Table 3b: Variants from GWAS of lung function


SNP	Chr	Locus	Risk Allele	Emphysema				Airway				Gas Trapping	
				%LAA-950		Perc15		Pi10		Wall Area Percent			
				All	Case	All	Case	All	Case	All	Case	All	Case
rs153916	5	<i>SPATA9-RHOBTB3</i>	T	0.001	0.02	2x10⁻⁵	0.02	-0.2	-0.3	0.9	-0.7	0.002	0.1
rs1529672	3	<i>RARB</i>	C	8x10 ⁻⁴	0.06	2x10⁻⁴	0.08	0.5	-1	0.1	0.9	2x10 ⁻⁴	0.03
rs2284746	1	<i>MFAP2</i>	G	0.002	0.2	0.002	0.1	-0.06	-0.5	0.9	1	8x10 ⁻⁴	0.07
rs12899618	15	<i>THSD4</i>	A	0.003	0.2	0.02	0.3	0.7	0.4	0.02	0.3	0.003	0.6
rs7765379	6	<i>HLA-DQB1</i>	T	0.004	0.05	0.04	0.08	-0.4	-0.5	-0.4	-0.2	0.2	0.9
rs9978142	21	<i>KCNE2-LINC00310</i>	T	0.005	0.06	0.04	0.07	-0.01	-0.05	-0.5	-0.9	0.04	0.004

rs3817928	6	<i>GPR126</i>	A	0.01	0.5	0.01	0.8	-0.1	-0.3	0.4	0.4	0.006	0.2
rs1036429	12	<i>CCDC38</i>	C	0.04	0.03	0.01	0.06	-0.5	-0.5	0.1	0.5	0.04	0.4
rs11134779	5	<i>ADAM19</i>	G	0.02	0.1	0.01	0.2	0.5	0.3	0.5	-0.7	0.04	0.08
rs11172113	12	<i>LRP1</i>	T	0.04	-0.9	0.2	-0.6	0.4	0.6	0.5	0.09	9x10-5	0.2
rs993925	1	<i>TGFB2-LYPLAL1</i>	C	0.2	-0.3	0.1	-0.1	-0.8	-0.6	-1	-0.4	0.004	0.9
rs7594321	2	<i>DNER</i>	C	0.2	0.6	0.1	0.8	-0.4	0.3	-0.5	-1	0.07	0.2
rs2798641	6	<i>ARMC2</i>	T	0.5	0.3	0.6	-0.4	0.1	0.03	8x10-4	0.004	0.06	-0.7
rs10516526	4	<i>GSTCD/INTS12/NPNT</i>	A	0.4	-0.3	0.4	-0.2	0.04	0.009	0.001	0.003	0.006	0.3
rs11168048	5	<i>HTR4</i>	T	0.05	0.5	0.09	0.8	0.06	0.2	0.002	0.07	0.3	-0.5
rs2865531	16	<i>CFDP1</i>	A	-1	-0.7	-0.9	-0.8	0.08	0.4	0.007	0.07	0.3	-0.3
rs2571445	2	<i>TNS1</i>	A	0.4	0.2	-0.3	0.4	1	-0.5	0.008	0.1	-0.2	-0.7
rs11654749	17	<i>KCNJ2</i>	T	-0.1	-0.05	-0.09	-0.04	0.4	-0.5	0.02	1	-0.5	-0.3
rs1344555	3	<i>MECOM</i>	T	-0.8	-1	-0.5	-0.8	0.5	0.7	0.3	0.05	-0.1	0.9
rs2857595	6	<i>NCR3-AIF1</i>	A	0.9	0.6	0.7	0.3	0.3	0.09	0.3	0.06	-0.6	0.6
rs11001819	10	<i>C10orf11</i>	G	-0.04	-0.01	-0.02	-0	0.7	0.8	0.07	0.1	-1	-0.1
rs16909898	9	<i>PTCH1</i>	G	-1	-0.1	0.7	-0.2	0.5	-0.8	0.2	-0.9	0.1	-0.9
rs12447804	16	<i>MMP15</i>	T	-0.2	-0.3	-0.3	-0.3	0.6	0.5	0.7	0.2	-0.6	-0.6
rs7068966	10	<i>CDC123</i>	C	-0.5	-0.5	0.8	-1	-0.1	-0.1	0.2	0.9	0.8	-0.7
rs6903823	6	<i>ZKSCAN3</i>	G	-0.7	0.9	1	0.8	0.7	0.9	0.9	-0.9	-0.4	-0.7
rs12477314	2	<i>HDAC4-FLJ43879</i>	C	-0.3	-0.08	-0.3	-0.1	-0.01	-0.05	-0.6	-0.5	-0.7	-0.1

Figures

Figure 1: Local association plots for genome-wide significant loci. a-e) %LAA-950, f) wall area percent, g) % gas trapping.

Additional data are available in the Supplement.



Title

A Genome-wide Association Study of Emphysema and Airway Quantitative Imaging Phenotypes

Michael H. Cho^{1,2}, Peter J. Castaldi¹, Craig P. Hersh^{1,2}, Brian D. Hobbs^{1,2}, R. Graham Barr⁴, Ruth Tal-Singer⁵, Per Bakke⁶, Amund Gulsvik⁶, Raúl San José Estépar³, Edwin Van Beek⁷, Harvey O. Coxson⁸, David A. Lynch⁹, George R. Washko², Nan M. Laird¹⁰, James D. Crapo⁹, Terri H. Beaty¹¹, Edwin K. Silverman^{1,2}, on behalf of the NETT Genetics, ECLIPSE, and COPDGene Investigators

¹Channing Division of Network Medicine, ²Division of Pulmonary and Critical Care Medicine, and ³Laboratory of Mathematics in Imaging, Department of Radiology, Brigham and Women's Hospital, Boston, MA; ⁴Department of Medicine, College of Physicians and Surgeons, and Department of Epidemiology, Mailman School of Public Health, Columbia University, New York, New York, ⁵GlaxoSmithKline Research and Development, King Of Prussia, PA, ⁶Department of Clinical Science, University of Bergen, Norway, ⁷Queens Medical Research Institute, University of Edinburgh, Department of Radiology and Department of Biomedical Engineering, University of Iowa; ⁸Department of Radiology, University of British Columbia, Vancouver, Canada ⁹National Jewish Health, Denver, CO; ¹⁰Harvard School of Public Health, Boston, MA; ¹¹Johns Hopkins University Bloomberg School of Public Health, Baltimore, MD.

Supplemental Data

Supplemental Methods

Study Populations

Evaluation of COPD Longitudinally to Identify Predictive Surrogate End-points

(ECLIPSE; SCO104960, NCT00292552, www.eclipse-copd.com): ECLIPSE cases and controls were aged 40-75 with at least a 10 pack-year smoking history without other respiratory diseases and without known alpha-1 antitrypsin deficiency. Cases were GOLD Grade 2 and above (post-bronchodilator forced expiratory volume in 1 second (FEV_1) < 80% predicted and FEV_1 /forced vital capacity (FVC) < 0.7); controls had no evidence of obstruction and FEV_1 > 85% predicted. Details of the ECLIPSE study have been previously published(1).

Genotyping was performed using the Illumina HumanHap 550 V3 (Illumina, San Diego, CA), and BeadStudio quality control, including reclustering on project samples was performed following Illumina guidelines. Quality control was performed using Python (www.python.org) and R (www.r-project.org) scripts in conjunction with plink (v1.05). Subjects and markers with a call rate of < 95% were excluded. Population stratification exclusion and adjustment on self-reported white subjects was performed using EIGENSOFT Version 2.0. Details of the genotyping and previous genome-wide association have been published(2). Imputation was updated using MaCH and minimac with the 1000 Genomes Phase I v3 EUR reference panel as previously described(3), resulting in a total of 11,040,911 variants with $R_{sq} > 0.3$.

Low-dose (120kVp and 40mAs) CT scans were performed at baseline, 1 year, and 3 year time points; baseline scans were used for the current analysis. All scans were performed using multidetector CT scans (GE Healthcare, Milwaukee, Wis. or Siemens Healthcare, Erlangen,

Germany) and images were reconstructed using 1.0mm (Siemens) or 1.25mm (GE) contiguous slices and an intermediate spatial frequency reconstruction algorithm. CT scanners were calibrated regularly using standard water calibration phantoms. All CT scans were analyzed at the University of British Columbia using Pulmonary Workstation 2.0 software (VIDA Diagnostics, Coralville, IA, U.S.A.). Airways were segmented using a region growing algorithm using the third (segmental) to fifth generation airways(4, 5). Wall area percent was calculated using the mean value of measurements for selected segmental airways (the same as used for COPDGene below) across all lobes.

National Emphysema Treatment Trial (NETT, www.nhlbi.nih.gov/health/prof/lung/nett/): NETT subjects had severe airflow obstruction by post-bronchodilator spirometry ($FEV_1 < 45\%$ predicted) and evidence of emphysema on computed tomography (CT). Subjects with significant sputum production or bronchiectasis were excluded. Details of the NETT trial have been published(6).

For the NETT Genetics Ancillary Study, we genotyped a subset of 382 self-reported white subjects without severe alpha-1 antitrypsin deficiency with available blood for genotyping who provided written consent. Genotyping was performed using the Illumina Quad 610 array (Illumina, San Diego, CA), with quality control, population stratification adjustment, and imputation procedures as previously described previously(2). A separate set of principal components was calculated for the NETT cases. Imputation was updated using MaCH and minimac with the 1000 Genomes Phase I v3 EUR reference panel(3) , resulting in a total of 10,659,967 variants with $R_{sq} > 0.3$.

NETT CT scans were performed on one of three types of scanners (General Electric, Fairfield, CT; Siemens, Malvern, PA; or Picker International, Toronto, ON, Canada) with a range of 2- to 8-mm slice thickness, with 75% of the scan data from 4 to 5 mm. Densitometric measures were performed with the Pulmonary Analysis Software Suite (PASS, Iowa City, IA). Airway measurements were obtained using 3D Slicer (www.Slicer.org) and Airway Inspector (www.airwayinspector.org) at Brigham and Women's Hospital. The full width at half-maximum (FWHM) method was used to measure the wall thickness and wall area of each airway.

Norway (GenKOLS, Genetics of Chronic Obstructive Lung Disease, GSK code RES11080):

GenKOLS cases and controls had at least a > 2.5 pack year smoking history. Cases had post-bronchodilator FEV1 < 80% predicted and FEV1/FVC < 0.7, while controls had normal spirometry. Subjects with severe alpha-1 antitrypsin deficiency and other lung diseases (aside from asthma) were excluded. Details of the GenKOLS study have been previously published(7).

Genotyping was performed using Illumina HumanHap 550 arrays (Illumina, San Diego, CA), with quality control, population stratification adjustment, and imputation procedures as described previously. A separate set of principal components was calculated for the subset of subjects with CT imaging data. Imputation was updated using MaCH and minimac with the 1000 Genomes Phase I v3 EUR reference panel(3), resulting in a total of 10,657,975 variants with Rsq > 0.3.

High-resolution CT chest scans were performed on a subset of the cohort using a GE LightSpeed Ultra. A low spatial frequency reconstruction algorithm was used for density measurements, and a high spatial frequency algorithm (bone) for airway measurements. Images were analyzed at the James Hogg iCAPTURE Centre (Vancouver, BC, Canada). Emphysema extent was assessed on lung images segmented using a modified boarder tracing algorithm with prior position

knowledge, and the extent of emphysema was assessed using the percentage of lung voxels with attenuation values less than -950 Hounsfield units (HU). Airways with an internal perimeter > 6mm were identified on the CT scans and measured using the Full Width at Half Maximum algorithm. Details on the imaging techniques in GenKOLS have been previously described(8).

COPDGene (NCT00608764, www.copdgene.org). COPDGene subjects were of non-Hispanic white or African-American ancestry, aged 45-80 years old, with a minimum of 10 pack-years of smoking, and without a history of lung disease other than asthma. Subjects found to have evidence of other lung disease on CT, such as significant bronchiectasis or interstitial lung disease, were excluded from the current analysis. Genotyping was performed by Illumina (San Diego, CA) on the HumanOmniExpress array, with quality control and imputation as previously described(3), resulting in a total of 11,437,352 variants for non-Hispanic whites and 22,904,273 for African-Americans with $R_{sq} > 0.3$.

CT chest imaging was performed on all subjects using a standardized protocol(9). Quantitative analysis utilized the lower-spatial-resolution smooth reconstruction algorithm. Analysis of emphysema severity was performed on segmented lung images by using the Slicer software package (<http://www.slicer.org/>). Emphysema percentage was defined as all lung voxels with a CT attenuation value of less than -950 HU. Airway analysis was performed by using the VIDA Pulmonary Workstation, version 2.0 (Vida Diagnostics, Coralville, Iowa, <http://www.vidadiagnostics.com/>). Measurements were obtained along the center line of the lumen, in the middle third of the airway segment, for one segmental airway of each lung lobe including the lingula; the mean value across all lobes was used for analysis. Details of the imaging techniques have been described previously(10).

Additional Genetic Analysis Methods

Imputed genotypes were included for analysis if they had an R^2 of 0.3 or greater. Individual genetic variants were included in the meta-analysis if they were missing in no more than one study (except for gas trapping, where the variant was required to be present in both COPDGene populations); variants with minor allele frequency $< 1\%$ overall or $< 0.5\%$ in individual studies were excluded, resulting in 6.9 (gas trapping) to 7.6 million (all other phenotypes) total analyzed variants. All variants were oriented to the '+' strand of the hg19 reference assembly. P-values were not adjusted for multiple comparisons.

Our primary analyses were performed in all subjects, with a method used to specifically address ascertainment. We additionally assessed the impact of each of the top variants in cases and non-cases separately using the same methods as for the overall meta-analysis. For results in the *SERPINA10* locus, we performed a meta-analysis conditioning on the *SERPINA1* Z allele by performing a linear regression including this SNP as a covariate in the model, and performing a meta-analysis on the target SNP.

To determine whether loci previously described in association with lung function were enriched for nominally significant ($P < 0.05$) associations in our quantitative imaging, we performed a Fisher's exact test. To determine whether any of the variants that we identified in this analysis were expression quantitative trait loci in lung, we searched the published dataset of Hao et al (11) and data from the GTEx consortium. Since Hao et al report only significant genotyped loci, we searched for variants in linkage disequilibrium with our top-reported variants using plink. VEGAS version 0.8.27 (12) gene-based analysis was performed using the CEU reference

haplotypes and including the top 20 percent of SNPs for a given gene. For the GRAIL
(<http://www.broadinstitute.org/mpg/grail/grail.php>)(13) analysis, HapMap2 CEU variants were
pruned for linkage disequilibrium using plink(14) with 250kb windows and an r^2 of 0.1. Results
with overall P-value $< 1 \times 10^{-4}$ were input as seed and query regions, including text from PubMed
articles up to May 2012. For the analysis using iGSEA4GWAS(15), default settings of 500kb up
and downstream boundaries and canonical pathways was used. For DEPICT(16)
(<http://www.broadinstitute.org/mpg/depict/>), SNPs were pruned to 500kb boundaries with an r^2
of 0.05. For INRICH(17), input files were pruned using an r^2 of 0.05 using a range of 20kb up
and downstream with 10,000 replicates. MAGENTA(18) was run using version July 2011,
under default settings. Overlap between results from these analyses was examined using an FDR
 < 0.05 for iGSEA4GWAS, $P < 0.005$ for DEPICT, $P < 0.05$ for INRICH, and nominal GSEA
75th percentile $P < 0.05$, to allow similar number of results in each dataset.

For the analysis of enhancer and promoter enrichment in ENCODE data, we used Haploreg
v2(19), using SNPs with GWAS P-values of $< 1 \times 10^{-6}$ for the top GWAS results, an r^2 of 0.8 and
using 1000 Genomes EUR Pilot data as background for enrichment. Briefly, Haploreg calculates
enrichment using the background set of variants to determine the level of overlap of specifically
annotated regions from the ENCODE project, and calculates an uncorrected binomial P-value.

Linkage disequilibrium between SNPs was estimated using the 1000 Genomes reference data in
SNAP(20), the 1000 Genomes EUR reference data, or (for the calculation with the reported
DLCI variant) the imputed genotypes in the African-American COPDGene samples, and
calculated using plink. All chromosomal positions are given using the NCBI37/hg19 assembly,
and alleles are referenced to the + strand.

152 **Supplemental Results**

153 **Genome-wide Association Quality Control**

154 None of the individual genome-wide association results for each cohort and phenotype
155 demonstrated evidence of substantial inflation of p-values (λ_{GC} range 1.0 – 1.02). For the meta-
156 analyses, the fixed effects analysis for Pi10 in all subjects demonstrated minimal evidence of
157 inflation ($\lambda_{GC}=1.06$, $\lambda_{GC1000}=1.01$), the remainder of both fixed and modified random effects
158 studies did not show evidence of inflation ($\lambda_{GC} = 1.02$).

159

160 **Supplemental Tables**

Table S1: Detailed results for the top genome-wide association results. Results given for each cohort. For the analyses involving all subjects, the second line shows the P-values from the SPREG(21) analysis (for COPDGene and Norway) or for cases only (ECLIPSE).

Phenotype	Cohort	Closest Gene	Marker Name	COPDGene non-Hispanic Whites			COPDGene African-Americans			ECLIPSE			NETT			Norway		
				Beta	SE	P-value	Beta	SE	P-value	Beta	SE	P-value	Beta	SE	P-value	Beta	SE	P-value
%LAA-950	All	HHIP	rs13141641	0.16	0.024	7.6x10 ⁻¹¹	0.11	0.056	0.059	0.11	0.038	0.0031	0.004	0.067	0.95	0.12	0.076	0.12
						5.5x10 ⁻⁹			0.082			0.12						0.15
		CHRNA3	rs55676755	-0.13	0.025	9.4x10 ⁻⁸	-0.094	0.045	0.037	-0.092	0.039	0.018	-0.02	0.066	0.76	-0.097	0.077	0.21
						3.8x10 ⁻⁶			0.1			0.015						0.47
		AGER	rs2070600	-0.35	0.058	1.6x10 ⁻⁹	-0.22	0.18	0.21	-0.22	0.1	0.029	0.26	0.14	0.065	-0.1	0.18	0.56
						1.1x10 ⁻⁸			0.2			0.042						0.41
		DLC1	rs75200691	0.16	0.037	2.6x10 ⁻⁵	0.18	0.063	0.0042	0.13	0.057	0.027	0.11	0.097	0.25	0.17	0.11	0.11
						3.5x10 ⁻⁵			0.0057			0.082						0.1
		SERPINA10	rs45505795	-0.28	0.074	1.7x10 ⁻⁴	-0.56	0.23	0.013	-0.39	0.1	0.00011	-0.064	0.16	0.7	-0.64	0.21	0.0024
						4.4x10 ⁻⁴			0.033			0.0011						0.0069
Perc15, HU	All	DLC1	rs74834049	-3.3	0.7	3.0x10 ⁻⁶	-3.6	1.4	0.011	-3	1.5	0.052	-3.5	2.1	0.095	-5.3	2.3	0.02
						3.8x10 ⁻⁶			0.015			0.15						0.017
		HHIP	rs13141641	-2.5	0.45	1.7x10 ⁻⁸	-1	1.2	0.42	-3	0.99	0.0022	-0.25	1.4	0.86	-1.8	1.5	0.23
						9.3x10 ⁻⁷			0.52			0.041						0.3
WAP, %	All	MIR2054	rs142200419	1.3	0.27	1.1x10 ⁻⁶				1.8	0.67	0.0093	0.56	1.4	0.7	-2.8	0.71	6.9x10 ⁻⁵
						8.0x10 ⁻⁶						0.0016						7.7x10 ⁻⁵
Gas trapping, %	All	AGER	rs2070600	-0.24	0.042	1.4x10 ⁻⁸	-0.13	0.15	0.39									
						2.0x10 ⁻⁸			0.2									
		LINC00310	rs55706246	0.11	0.03	2.3x10 ⁻⁴	0.45	0.099	4.7x10 ⁻⁶									
						1.0x10 ⁻⁴			3.2x10 ⁻⁷									

Table S2: Additional results from each genome-wide study. Results with $P < 1 \times 10^{-6}$ in either the modified random effects or fixed effects analysis are shown.

Phenotype	Group	Chr	Marker Name	Closest Gene	Effect Allele	Allele Frequency		Modified Random Effects			Fixed Effects		
<i>Emphysema</i>						Nhw	Aa	P value	Beta	SE	P value	Nhw	Aa
%LAA-950, %	All	9	rs3919995	<i>ZNF462</i>	A	0.59	0.5	1.3×10^{-7}	-0.081	0.023	8.1×10^{-8}	-0.088	0.016
		20	rs183345681	<i>CHRNA4</i>	A	0.23	0.18	1.8×10^{-7}	-0.12	0.023	1.1×10^{-7}	-0.12	0.023
		14	rs117167774	<i>LOC100506433</i>	T	0.013	0.013	1.8×10^{-7}	0.47	0.23	0.00013	0.33	0.086
		2	rs360488	<i>FAM84A</i>	A	0.23	0.082	3.7×10^{-7}	0.09	0.038	3.0×10^{-7}	0.11	0.021
		1	rs7512679	<i>TGFB2</i>	T	0.24	0.47	4.5×10^{-7}	0.092	0.018	2.9×10^{-7}	0.092	0.018
		8	rs7823498	<i>NRG1</i>	T	0.79	0.73	4.6×10^{-7}	-0.098	0.019	3.1×10^{-7}	-0.098	0.019
		11	rs7947523	<i>MIR4300</i>	C	0.68	0.44	4.9×10^{-7}	-0.086	0.048	0.00014	-0.064	0.017
		20	rs2070755	<i>PCK1</i>	C	0.49	0.4	5.3×10^{-7}	0.11	0.047	0.00041	0.058	0.016
		8	rs10109725	<i>CSMD1</i>	T	0.03	0.0069	6.3×10^{-7}	0.28	0.14	8.6×10^{-6}	0.25	0.055
		5	rs924633	<i>DNAH5</i>	A	0.95	0.92	9.2×10^{-7}	0.18	0.092	8.1×10^{-5}	0.14	0.036
		4	rs62343714	<i>LOC401164</i>	T	0.092	0.16	1.2×10^{-6}	0.12	0.036	8.8×10^{-7}	0.13	0.026
		19	rs7937	<i>MIA-RAB4B</i>	T	0.57	0.3	1.5×10^{-6}	-0.08	0.016	9.7×10^{-7}	-0.08	0.016
	Cases	11	rs608194	<i>MMP12</i>	T	0.18	0.33	1.4×10^{-7}	0.05	0.074	2.9×10^{-5}	0.11	0.027
		6	rs72971709	<i>GRIK2</i>	A	0.013	0.0029	2.6×10^{-7}	0.38	0.31	2.6×10^{-5}	0.44	0.1
		18	rs12605822	<i>ANKRD12</i>	A	0.13	0.11	3.6×10^{-7}	0.17	0.072	3.0×10^{-6}	0.15	0.031
		14	rs3811345	<i>LINC00617</i>	A	0.87	0.86	4.4×10^{-7}	0.16	0.03	2.8×10^{-7}	0.16	0.03
		15	rs9788721	<i>AGPHD1</i>	T	0.62	0.62	5.5×10^{-7}	-0.1	0.025	3.5×10^{-7}	-0.11	0.021
		1	rs72482608	<i>PRRX1</i>	A	0.62	0.52	7.6×10^{-7}	-0.11	0.021	4.8×10^{-7}	-0.11	0.021
		5	rs13184316	<i>ARL15</i>	A	0.23	0.05	8.2×10^{-7}	0.07	0.1	0.78	-0.0073	0.027
Perc15, HU	All	1	rs72637224	<i>XCL2</i>	T	0.05	0.14	3.3×10^{-7}	3.6	1.2	2.1×10^{-7}	3.5	0.68
		16	rs9933712	<i>ERCC4</i>	A	0.021	0.38	4.2×10^{-7}	5.2	1.8	2.6×10^{-7}	3.7	0.72
		20	rs183345681	<i>CHRNA4</i>	A	0.23	0.18	4.7×10^{-7}	2.4	0.47	3.0×10^{-7}	2.4	0.47
		12	rs75751297	<i>FLJ31485</i>	A	0.47	0.36	6.6×10^{-7}	2.4	0.48	4.2×10^{-7}	2.4	0.48
		11	rs7125940	<i>MIR4300</i>	T	0.34	0.58	6.9×10^{-7}	-1.9	1	8.3×10^{-5}	-1.4	0.35
		15	rs144442299	<i>UNC13C</i>	T	0.018	0.0051	7.8×10^{-7}	-5.4	2.9	5.4×10^{-7}	-7.4	1.5
		20	rs2070755	<i>PCK1</i>	C	0.49	0.4	8.5×10^{-7}	-2.5	1.1	0.0092	-0.88	0.34

		3	rs111646341	LSAMP	A	0.97	0.98	9.0x10 ⁻⁷	5.8	1.8	5.7x10 ⁻⁷	5.6	1.1
		14	rs45505795	SERPINA10	C	0.038	0.0076	9.5x10 ⁻⁷	6.4	2.7	2.6x10 ⁻⁶	5.2	1.1
		4	rs10016562	TRPC3	T	0.62	0.73	1.0x10 ⁻⁶	1.6	0.5	6.4x10 ⁻⁷	1.7	0.35
		8	rs7823498	NRG1	T	0.79	0.73	1.0x10 ⁻⁶	2	0.4	6.4x10 ⁻⁷	2	0.4
		15	rs9788721	AGPHD1	T	0.62	0.62	1.1x10 ⁻⁶	1.7	0.35	6.7x10 ⁻⁷	1.7	0.35
		6	rs2647050	HLA-DQB1	T	0.65	0.65	1.2x10 ⁻⁶	1.6	0.48	7.7x10 ⁻⁷	1.7	0.35
		20	rs6080212	KIF16B	A	0.16	0.15	1.4x10 ⁻⁶	-2.2	0.45	8.8x10 ⁻⁷	-2.2	0.45
	Cases	10	rs139326003	MBL2	A	0.12	0.089	1.6x10 ⁻⁷	4.2	0.95	1.2x10 ⁻⁷	3.9	0.74
		11	rs185888204	OR8B3	A	0.11	0.11	1.9x10 ⁻⁷	-7.1	3	2.5x10 ⁻⁶	-6.1	1.3
		15	rs503464	CHRNA5	A	0.22	0.27	2.5x10 ⁻⁷	-3.2	0.6	1.5x10 ⁻⁷	-3.2	0.6
		18	rs12605822	ANKRD12	A	0.13	0.11	4.0x10 ⁻⁷	-3.3	1.4	5.3x10 ⁻⁷	-3.6	0.71
		1	rs72482608	PRRX1	A	0.62	0.52	5.0x10 ⁻⁷	2.5	0.48	3.2x10 ⁻⁷	2.5	0.48
		11	rs654600	MMP12	A	0.83	0.72	5.2x10 ⁻⁷	-1.7	1.6	5.1x10 ⁻⁵	-2.5	0.63
		4	rs13140744	TRPC3	T	0.38	0.26	8.9x10 ⁻⁷	-2.2	0.64	5.7x10 ⁻⁷	-2.4	0.48
		1	rs75565482	XCL2	A	0.95	0.91	1.1x10 ⁻⁶	5.1	1.6	7.1x10 ⁻⁷	5.2	1.1
		14	rs3811345	LINC00617	A	0.87	0.86	1.5x10 ⁻⁶	-3.4	0.7	9.5x10 ⁻⁷	-3.4	0.7
Airway													
Pi10	All	8	rs13281609	CSMD3	T	0.047	0.0079	3.2x10 ⁻⁷	-0.044	0.01	2.2x10 ⁻⁷	-0.043	0.0082
		11	rs113835537	CTSF	A	0.84	0.83	8.5x10 ⁻⁷	0.012	0.0023	5.4x10 ⁻⁷	0.012	0.0023
		1	rs654950	HIVEP3	C	0.42	0.12	8.6x10 ⁻⁷	-0.011	0.0055	3.5x10 ⁻⁶	-0.0089	0.0019
	Cases	3	rs168302	GRM7	T	0.66	0.87	9.8x10 ⁻⁸	-0.016	0.004	6.0x10 ⁻⁸	-0.017	0.0032
		9	rs4877691	FAM75D1	A	0.24	0.38	6.6x10 ⁻⁷	-0.017	0.0078	2.0x10 ⁻⁶	-0.016	0.0034
		2	rs115089939	LOC647012	T	0.99	1	1.1x10 ⁻⁶	-0.086	0.017	7.2x10 ⁻⁷	-0.086	0.017
		5	rs79581221	ATG10	T	0.014	0.0017	1.1x10 ⁻⁶	-0.077	0.016	7.4x10 ⁻⁷	-0.077	0.016
WAP	All	1	rs12724666	PDZK1P1	A	0.033	0.0092	8.7x10 ⁻⁸	1.1	0.2	5.9x10 ⁻⁸	1.1	0.2
		8	rs2513900	AZIN1	C	0.51	0.74	2.6x10 ⁻⁷	0.23	0.043	1.7x10 ⁻⁷	0.23	0.043
		17	rs3826538	RPA1	T	0.072	0.27	1.5x10 ⁻⁶	-0.35	0.071	9.3x10 ⁻⁷	-0.35	0.071
	Cases	3	rs76493322	GRM7	A	0.46	0.45	3.1x10 ⁻⁷	-0.36	0.069	2.0x10 ⁻⁷	-0.36	0.069
		2	rs10932600	ATIC	A	0.62	0.73	1.3x10 ⁻⁶	-0.32	0.065	8.4x10 ⁻⁷	-0.32	0.065
		1	rs61797053	KIAA1324	A	0.067	0.019	1.5x10 ⁻⁶	0.67	0.14	9.5x10 ⁻⁷	0.67	0.14
Gas Trapping													

All	4	rs1512281	<i>HHIP-AS1</i>	A	0.59	0.88	2.3x10 ⁻⁷	0.082	0.016	1.9x10 ⁻⁷	0.082	0.016
	8	rs74834049	<i>DLC1</i>	A	0.11	0.082	6.1x10 ⁻⁷	0.12	0.024	5.0x10 ⁻⁷	0.12	0.024
	1	rs6669119	<i>PAX7</i>	T	0.1	0.12	9.90E-07	-0.14	0.062	1.60E-06	-0.11	0.024
	8	rs2844036	<i>ANKRD46</i>	A	0.78	0.88	1.10E-06	-0.11	0.022	8.60E-07	-0.11	0.022
	10	rs655766	<i>BAMBI</i>	T	0.28	0.22	1.20E-06	0.08	0.016	9.90E-07	0.08	0.016
Cases	12	rs10875912	<i>MLL2</i>	T	0.66	0.67	8.30E-08	-0.091	0.017	7.10E-08	-0.091	0.017
	20	rs430086	<i>MACROD2</i>	A	0.98	0.86	2.50E-07	0.16	0.16	9.80E-06	0.19	0.044
	2	rs72822868	<i>SNAR-H</i>	T	0.91	0.98	5.20E-07	0.23	0.046	4.30E-07	0.23	0.046
	12	rs2460882	<i>SP1</i>	T	0.84	0.38	6.40E-07	0.11	0.022	5.30E-07	0.11	0.022
	11	rs1789001	<i>OR9G4</i>	A	0.57	0.43	6.80E-07	0.079	0.024	5.60E-07	0.084	0.017
	6	rs12527942	<i>MRPL14</i>	T	0.03	0.048	9.40E-07	0.32	0.34	0.033	0.1	0.047
	8	rs13259853	<i>CSMD1</i>	A	0.44	0.099	1.00E-06	-0.088	0.022	8.30E-07	-0.09	0.018
	17	rs12449664	<i>NTN1</i>	A	0.14	0.084	1.10E-06	0.15	0.03	9.10E-07	0.15	0.03

Table S3: Lookup of top quantitative CT association results in all subjects within separate analyses in COPD cases and non-cases.

Phenotype	Chr	Marker Name	Closest Gene	Effect Allele	Cases						Non-cases					
					Modified Random Effects			Fixed Effects			Modified Random Effects			Fixed Effects		
					P value	Beta	SE	P value	Beta	Se	P value	Beta	SE	P value	Beta	Se
%LAA-950	4	rs13141641	HHIP	T	4.4×10^{-5}	0.09	0.030	3.6×10^{-5}	0.09	0.021	2.0×10^{-2}	0.05	0.021	1.5×10^{-2}	0.05	0.021
	15	rs55676755	CHRNA3	C	3.2×10^{-6}	-0.08	0.034	3.1×10^{-6}	-0.09	0.021	5.1×10^{-1}	0.02	0.022	4.4×10^{-1}	0.02	0.022
	6	rs2070600	AGER	T	1.9×10^{-2}	-0.08	0.095	3.1×10^{-2}	-0.12	0.054	1.9×10^{-3}	-0.11	0.126	4.4×10^{-3}	-0.18	0.073
	8	rs75200691	DLC1	T	6.3×10^{-3}	0.09	0.032	4.4×10^{-3}	0.09	0.032	1.4×10^{-4}	0.12	0.032	9.4×10^{-5}	0.12	0.032
	14	rs45505795	SERPINA10	C	3.9×10^{-4}	-0.21	0.056	2.6×10^{-4}	-0.21	0.056	1.8×10^{-2}	-0.18	0.073	1.4×10^{-2}	-0.18	0.073
Perc 15	8	rs74834049	DLC1	A	7.5×10^{-4}	-2.6	0.74	5.1×10^{-4}	-2.6	0.74	2.7×10^{-4}	-0.27	0.81	1.9×10^{-4}	-2.4	0.064
	4	rs13141641	HHIP	T	8.3×10^{-5}	-2.0	0.49	5.5×10^{-5}	-2.0	0.49	2.7×10^{-1}	-0.53	0.43	2.2×10^{-1}	-0.52	0.43
Airway																
WAP	4	rs142200419	MIR2054	T	3.1×10^{-4}	0.30	1	3.9×10^{-2}	0.7	0.34	3.7×10^{-3}	-0.29	1.23	1.7×10^{-2}	0.71	0.30
Gas trapping																
%	6	rs2070600	AGER	T	5.2×10^{-3}	-0.15	0.083	4.4×10^{-3}	-0.13	0.047	5.1×10^{-4}	-0.18	0.05	4.3×10^{-4}	-0.18	0.050
	21	rs55706246	LINC00310	A	3.7×10^{-3}	0.15	0.099	3.2×10^{-3}	0.09	0.032	2.2×10^{-2}	0.19	0.159	3.9×10^{-2}	0.08	0.038

Table S4: Top overall quantitative CT loci not previously reported in case-control association analyses for moderate-to-severe and severe COPD in COPDGene, ECLIPSE, GenKOLS, and NETT/NAS(3)

Chr	Marker Name	Closest Gene	Effect Allele	Moderate-To-Severe COPD		Severe COPD	
				P-value	Beta	P-value	Beta
6	rs2070600	AGER	T	2.9×10^{-4}	-0.35	1.4×10^{-5}	-0.45
8	rs75200691	DLC1	T	0.35	0.05	0.21	0.08
8	rs74834049	DLC1	A	0.39	-0.04	0.20	-0.08
14	rs45505795	SERPINA10	C	3.4×10^{-5}	0.42	1.6×10^{-5}	0.51
4	rs142200419	MIR2054	T	0.25	0.19	0.47	0.14

21	rs55706246	<i>LINC00310</i>	A	6.5×10^{-3}	-0.16	1.2×10^{-2}	-0.18
----	------------	------------------	---	----------------------	-------	----------------------	-------

References

1. Vestbo J, Anderson W, Coxson HO, Crim C, Dawber F, Edwards L, Hagan G, Knobil K, Lomas DA, MacNee W, Silverman EK, Tal-Singer R. Evaluation of COPD Longitudinally to Identify Predictive Surrogate End-points (ECLIPSE). *Eur Respir J* 2008;31:869–73.
2. Cho MH, Boutaoui N, Klanderman BJ, Sylvia JS, Ziniti JP, Hersh CP, DeMeo DL, Hunninghake GM, Litonjua AA, Sparrow D, Lange C, Won S, Murphy JR, Beaty TH, Regan EA, Make BJ, Hokanson JE, Crapo JD, Kong X, Anderson WH, Tal-Singer R, Lomas DA, Bakke P, Gulsvik A, Pillai SG, Silverman EK. Variants in FAM13A are associated with chronic obstructive pulmonary disease. *Nat Genet*, 2010/02/23 ed. 2010;42:200–202.
3. Cho MH, McDonald M-LN, Zhou X, Mattheisen M, Castaldi PJ, Hersh CP, Demeo DL, Sylvia JS, Ziniti J, Laird NM, Lange C, Litonjua AA, Sparrow D, Casaburi R, Barr RG, Regan EA, Make BJ, Hokanson JE, Lutz S, Dudenkov TM, Farzadegan H, Hetmanski JB, Tal-Singer R, Lomas DA, Bakke P, Gulsvik A, Crapo JD, Silverman EK, Beaty TH. Risk loci for chronic obstructive pulmonary disease: a genome-wide association study and meta-analysis. *lancet Respir Med* 2014;2:214–25.
4. Coxson HO, Dirksen A, Edwards LD, Yates JC, Agusti A, Bakke P, Calverley PM, Celli B, Crim C, Duvoix A, Fauerbach PN, Lomas DA, Macnee W, Mayer RJ, Miller BE, Müller NL, Rennard SI, Silverman EK, Tal-Singer R, Wouters EF, Vestbo J. The presence and progression of emphysema in COPD as determined by CT scanning and biomarker expression: a prospective analysis from the ECLIPSE study. *lancet Respir Med* 2013;1:129–36.
5. Gietema HA, Müller NL, Fauerbach PVN, Sharma S, Edwards LD, Camp PG, Coxson HO. Quantifying the extent of emphysema: factors associated with radiologists' estimations and quantitative indices of emphysema severity using the ECLIPSE cohort. *Acad Radiol* 2011;18:661–71.
6. Fishman A, Martinez F, Naunheim K, Piantadosi S, Wise R, Ries A, Weinmann G, Wood DE. A randomized trial comparing lung-volume-reduction surgery with medical therapy for severe emphysema. *N Engl J Med* 2003;348:2059–73.
7. Pillai SG, Ge D, Zhu G, Kong X, Shianna K V, Need AC, Feng S, Hersh CP, Bakke P, Gulsvik A, Ruppert A, Lodrup Carlsen KC, Roses A, Anderson W, Rennard SI, Lomas DA, Silverman EK, Goldstein DB. A genome-wide association study in chronic obstructive pulmonary disease (COPD): identification of two major susceptibility loci. *PLoS Genet*, 2009/03/21 ed. 2009;5:e1000421.
8. Grydeland TB, Dirksen A, Coxson HO, Eagan TML, Thorsen E, Pillai SG, Sharma S, Eide GE, Gulsvik A, Bakke PS. Quantitative computed tomography measures of

- emphysema and airway wall thickness are related to respiratory symptoms. *Am J Respir Crit Care Med* 2010;181:353–9.
9. Regan EA, Hokanson JE, Murphy JR, Make B, Lynch DA, Beaty TH, Curran-Everett D, Silverman EK, Crapo JD. Genetic epidemiology of COPD (COPDGene) study design. *COPD*, 2010/03/11 ed. 7:32–43.
 10. Han MK, Kazerooni EA, Lynch DA, Liu LX, Murray S, Curtis JL, Criner GJ, Kim V, Bowler RP, Hanania NA, Anzueto AR, Make BJ, Hokanson JE, Crapo JD, Silverman EK, Martinez FJ, Washko GR, The COPDGene Investigators F. Chronic obstructive pulmonary disease exacerbations in the COPDGene study: associated radiologic phenotypes. *Radiology* 2011;261:274–82.
 11. Hao K, Bossé Y, Nickle DC, Paré PD, Postma DS, Laviolette M, Sandford A, Hackett TL, Daley D, Hogg JC, Elliott WM, Couture C, Lamontagne M, Brandsma C-A, van den Berge M, Koppelman G, Reicin AS, Nicholson DW, Malkov V, Derry JM, Suver C, Tsou JA, Kulkarni A, Zhang C, Vessey R, Opitck GJ, Curtis SP, Timens W, Sin DD. Lung eQTLs to help reveal the molecular underpinnings of asthma. *PLoS Genet* 2012;8:e1003029.
 12. Liu JZ, Mcrae AF, Nyholt DR, Medland SE, Wray NR, Brown KM, Investigators A, Hayward NK, Montgomery GW, Visscher PM. A Versatile Gene-Based Test for Genome-wide Association Studies. *Am J Hum Genet* 2010;87:139–45.
 13. Raychaudhuri S, Plenge RM, Rossin EJ, Ng ACY, Purcell SM, Sklar P, Scolnick EM, Xavier RJ, Altshuler D, Daly MJ. Identifying relationships among genomic disease regions: predicting genes at pathogenic SNP associations and rare deletions. *PLoS Genet* 2009;5:e1000534.
 14. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J, Sklar P, De Bakker PIW, Daly MJ, others. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* 2007;81:559–75.
 15. Zhang K, Cui S, Chang S, Zhang L, Wang J. i-GSEA4GWAS: a web server for identification of pathways/gene sets associated with traits by applying an improved gene set enrichment analysis to genome-wide association study. *Nucleic Acids Res*, 2010/05/04 ed. 2010;38:W90–5.
 16. Wood AR, Esko T, Yang J, Vedantam S, Pers TH, Gustafsson S, Chu AY, Estrada K, Luan J, Kutalik Z, Amin N, Buchkovich ML, Croteau-Chonka DC, Day FR, Duan Y, Fall T, Fehrmann R, Ferreira T, Jackson AU, Karjalainen J, Lo KS, Locke AE, Mägi R, Mihailov E, Porcu E, Randall JC, Scherag A, Vinkhuyzen AAE, Westra H-J, *et al*. Defining the role of common variation in the genomic and biological architecture of adult human height. *Nat Genet* 2014;advance on:1173–86.

17. Lee PH, O'Dushlaine C, Thomas B, Purcell SM. INRICH: interval-based enrichment analysis for genome-wide association studies. *Bioinformatics* 2012;28:1797–9.
18. Segre A V, Groop L, Mootha VK, Daly MJ, Altshuler D. Common inherited variation in mitochondrial genes is not enriched for associations with type 2 diabetes or related glycemic traits. *PLoS Genet*, 2010/08/18 ed. 2010;6.:
19. Ward LD, Kellis M. HaploReg: a resource for exploring chromatin states, conservation, and regulatory motif alterations within sets of genetically linked variants. *Nucleic Acids Res* 2012;40:D930–4.
20. Johnson AD, Handsaker RE, Pulit SL, Nizzari MM, O'Donnell CJ, de Bakker PIW. SNAP: a web-based tool for identification and annotation of proxy SNPs using HapMap. *Bioinformatics* 2008;24:2938–9.
21. Lin DY, Zeng D. Proper analysis of secondary phenotype data in case-control association studies. *Genet Epidemiol* 2009;33:256–65.